

# Chapter 4

## Definition of Artificial Neural Network

### ABSTRACT

*In living creatures, the brain is the control unit and it can be divided in different anatomic and functional sub-units. An artificial neural network is a computational system for processing information as a response to external stimuli, which consists of a set of highly interconnected processing elements called neurons. It is very useful to have some knowledge of the way the biological nervous system is organized, since the artificial neural network is an inspiration of the biological neural networks. This chapter is an explanation of the Artificial Neural Network (ANN). The biological and mathematical definition of a neural network is provided and the activation functions effective for processing are listed. Some figures are collected for better understanding.*

### 1 BIOLOGICAL NEURAL NETWORK

In the nervous system of the living creatures, there are fluid-filled sacs which bound by a lipid bilayer for separating the intracellular contents from the extracellular space and they are called Neurons, or brain cells. Inside the body, neurons are responsible to maintain a negative internal voltage, which is related to the extracellular space; ion channels and pumps maintain this potential difference. In most neurons of the central nervous system, spike is responsible to send the signals of neural activity, or rapid intracellular depolarization followed by repolarization; in order to adjust

DOI: 10.4018/978-1-4666-6146-2.ch004

### ***Definition of Artificial Neural Network***

the neurons, it is necessary to communicate information about a neuron's activity. Some neurons communicate with simple resistive coupling, via channels that allow direction flow. However, for higher animals, most neurons in the central nervous system (CNS), communicate through chemical synapses: triggering the release of chemicals using the neural spike is called neurotransmitters into the extracellular space. These neurotransmitters bind to ion channels in adjacent neurons, causing a brief ionic current to flow into the neuron. The resulting current flow in the recipient neuron will be depolarizing, or hyperpolarizing and it depends on whether the neurotransmitter is excitatory or inhibitory, respectively.

It is very useful to have some knowledge of the way the biological nervous system is organized, since the artificial neural network is an inspiration of the biological neural networks.

There is difference among the nervous system of creatures. Most living creatures, which have the ability to adapt to a changing environment, need a controlling unit, which is able to learn. Higher developed animals and humans use very complex networks of highly specialized neurons to perform this task.

In the living creatures, the brain is the control unit and it can be divided in different anatomic and functional sub-units. Each unit is responsible to do certain tasks like vision, hearing, motor and sensor control. The brain is connected by nerves to the sensors and actors in the rest of the body.

There are a large number of neurons, about  $10^{11}$  in average. This can be considered as the basic building bricks for the central nervous system (CNS). The connection points of the neurons are called synapses. The complexity of the brain is because of the massive number of highly interconnected simple units working in parallel, with an individual neuron receiving input from up to 10000 others (Bishop, 1995).

The structure and its processes in a simple cell are enormously complex. Even the 'dendrites' that originating from the cell body are thin and widely branching fibers, reaching out in different directions and make connections to a larger number of cells within the cluster. Axons make the connections from the other cells to dendrites of one cell or directly to the body of the cell. There is only one axon per neuron. It is a single and long fiber, which transports the output signal of the cell as electrical impulses (action potential) along its length. The end of the axon may divide in many branches, which are then connected to other cells. The branches have the function to fan out the signal to many other inputs (Hassoun, 1995).

The structure of neuron has four main regions. There are two offshoots from the cell body, the dendrites, and the axon, which end in presynaptic terminals. The heart of the cell is called the cell body that contains the nucleus and maintaining protein synthesis. A neuron has many dendrites in a treelike structure, which branch out, and receive signals from other neurons. Each neuron has one axon that grows out from a part of the cell body called the axon hillock. The axon hillock generates

12 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: [www.igi-global.com/chapter/definition-of-artificial-neural-network/110999](http://www.igi-global.com/chapter/definition-of-artificial-neural-network/110999)

## Related Content

---

### Optimization of Evolutionary Algorithm Using Machine Learning Techniques for Pattern Mining in Transactional Database

Logeswaran K., Suresh P., Savitha S. and Prasanna Kumar K. R. (2020). *Handbook of Research on Applications and Implementations of Machine Learning Techniques* (pp. 173-200).  
[www.irma-international.org/chapter/optimization-of-evolutionary-algorithm-using-machine-learning-techniques-for-pattern-mining-in-transactional-database/234124](http://www.irma-international.org/chapter/optimization-of-evolutionary-algorithm-using-machine-learning-techniques-for-pattern-mining-in-transactional-database/234124)

### Energy of Graphs

Harishchandra S. Ramane (2020). *Handbook of Research on Advanced Applications of Graph Theory in Modern Society* (pp. 267-296).  
[www.irma-international.org/chapter/energy-of-graphs/235540](http://www.irma-international.org/chapter/energy-of-graphs/235540)

### Trigonometric Polynomial Higher Order Neural Network Group Models and Weighted Kernel Models for Financial Data Simulation and Prediction

Lei Zhang, Simeon J. Simoff and Jing Chun Zhang (2009). *Artificial Higher Order Neural Networks for Economics and Business* (pp. 484-503).  
[www.irma-international.org/chapter/trigonometric-polynomial-higher-order-neural/5296](http://www.irma-international.org/chapter/trigonometric-polynomial-higher-order-neural/5296)

### Complex-Valued Neural Networks for Equalization of Communication Channels

Rajoo Pandey (2009). *Complex-Valued Neural Networks: Utilizing High-Dimensional Parameters* (pp. 168-193).  
[www.irma-international.org/chapter/complex-valued-neural-networks-equalization/6769](http://www.irma-international.org/chapter/complex-valued-neural-networks-equalization/6769)

### Generative Adversarial Neural Networking of Agents: Avatars as Tools for Financial Modelling

Vladimir Soloviev, Vsevolod Chernyshenko, Vadim Feklin, Ekaterina Zolotareva and Nikita Titov (2020). *Avatar-Based Control, Estimation, Communications, and Development of Neuron Multi-Functional Technology Platforms* (pp. 107-120).  
[www.irma-international.org/chapter/generative-adversarial-neural-networking-of-agents/244788](http://www.irma-international.org/chapter/generative-adversarial-neural-networking-of-agents/244788)