

Chapter 8

Rift Valley Fever and the Changing Environment: A Case Study in East Africa

Johanna Lindahl

*Swedish University of Agricultural
Sciences, Sweden & International
Livestock Research Institute, Kenya*

Bernard Bett

*International Livestock Research
Institute, Kenya*

Timothy Robinson

*International Livestock Research
Institute, Kenya*

Delia Grace

*International Livestock Research
Institute, Kenya*

ABSTRACT

Rift Valley fever is a severe disease affecting both humans and animals. The Rift Valley fever virus can be transmitted by body fluids, and the most common way for humans to get infected is from animals. The virus is also vector-borne and can be transmitted by many species of mosquitoes. As with other vector-borne diseases, the epidemiology may vary in response to environmental changes. Here the effects of climate and land use changes on Rift Valley fever, as well as on other vector-borne diseases, are discussed. The effect of irrigation in East Africa on inter-epidemic transmission of RVF is discussed in greater detail, followed by recommendations for future research and actions.

DOI: 10.4018/978-1-5225-0553-2.ch008

INTRODUCTION

The last century has seen a period of ecological change, unprecedented in recent times, with dramatic reductions in pristine habitats, ecosystem services and biodiversity along with equally dramatic increases in numbers of people and of domestic animals. Most land use practices change the environment and are due to human influence, often related to feeding the growing population, such as increased irrigation and land deforestation to allow crop cultivation. Environmental change is ubiquitous, and new diseases continue to emerge, while concerns about the impact of environmental change on human health grow. However, while it seems intuitive that degradation of ecosystems will increase the risk of disease, detailed knowledge on how specific environmental changes in a given context affect disease transmission is often lacking. As a consequence, decision makers may inadvertently increase health risks.

This chapter uses the example of Rift Valley fever (RVF) to explore relations between disease dynamics and environmental change. Two main drivers, climate and land use changes, can be identified. We focus on irrigation as a type of land use change. In addition, RVF is characterized, describing the knowns and unknowns about its epidemiology and recapitulating the history and impact of RVF outbreaks in east Africa. Next we discuss relations between RVF and climate and land use change. Findings from an ongoing study, which shows that an irrigated area can support endemic transmission of RVF, without any signs of outbreaks, are also presented and the relevance of this is discussed.

IMPORTANCE AND DRIVERS OF EMERGING DISEASE

Emerging infectious diseases in both animals and humans cause major economic and health burdens in every part of the world. On average, a new human disease appears every four months and around 75% of emerging diseases are zoonotic (Jones et al., 2008). Most originate from wildlife, and the study of disease emergence has a strong focus on wildlife. However, economically important emerging diseases often involve domestic animals. For example, between 1997 and 2009, six major emerging diseases have together cost at least 80 billion USD: the Nipah virus outbreak in Malaysia, West Nile fever in the USA, severe acute respiratory syndrome (SARS, starting in Asia), highly pathogenic avian influenza (HPAI, starting in Asia), bovine spongiform encephalopathy (BSE, starting in the UK) and RVF in East Africa (World Bank, 2012). In all of these, livestock or animals farmed for human consumption provided either a reservoir or a bridge to transmit the disease to people. Later outbreaks of emerging infectious diseases, such as Middle East respiratory syndrome (MERS) and the Ebola outbreak in West Africa, were also caused by viruses with

25 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/chapter/rift-valley-fever-and-the-changing-environment/162357

Related Content

Nutraceutical and Functional Foods in Cancer Management and Therapy

Sakshi Bajaj and Satish Manchanda (2019). *Nutraceutical and Functional Foods in Disease Prevention* (pp. 341-390).

www.irma-international.org/chapter/nutraceutical-and-functional-foods-in-cancer-management-and-therapy/207983

Business Executives' Perceptions of Responsible Leadership and Corporate Social Responsibility for Stakeholders' Health and Wellbeing

Frida Stål, Anneli Marttila and Gloria Macassa (2022). *International Journal of Applied Research on Public Health Management* (pp. 1-14).

www.irma-international.org/article/business-executives-perceptions-responsible-leadership/290378

Pharmacy Technology to Better Public Health: An Exploration of New Models of Supply and Use of Technology – A Regional United Kingdom Quantitative Study

Shahid Muhammad, Hooman Safaei and Tariq Muhammad (2019). *International Journal of Applied Research on Public Health Management* (pp. 1-20).

www.irma-international.org/article/pharmacy-technology-to-better-public-health/218865

Towards a Comprehensive National Health System for the United States

David Stewart, Sheila R. Goodwin and Nikolaos Karagiannis (2023). *Assessing the Need for a Comprehensive National Health System in the United States* (pp. 41-56).

www.irma-international.org/chapter/towards-a-comprehensive-national-health-system-for-the-united-states/315879

Twitter and Its Role in Health Information Dissemination: Analysis of the Micro-Blog Posts of Health-Related Organisations

Dan Dumbrell and Robert Steele (2017). *Public Health and Welfare: Concepts, Methodologies, Tools, and Applications* (pp. 372-388).

www.irma-international.org/chapter/twitter-and-its-role-in-health-information-dissemination/165821