Chapter 6 Nutraceutical Intervention of Phytosterols in Cardiovascular Aging

Charu Gupta Amity University, India

Dhan Prakash Amity University, India

Sneh Gupta RGPG College, India

ABSTRACT

Phytosterols and phytostanols are a large group of compounds that are found exclusively in plants. They are structurally and functionally related to cholesterol but differ from cholesterol in the structure of the side chain. Phytosterols lower total and low-density lipoprotein (LDL) blood cholesterol by preventing cholesterol absorption from the intestine, so they have been known as blood cholesterol-lowering agents. Phytosterols are naturally found in fruits, vegetables, nuts, and mainly oils. Dietary phytosterol intakes normally range from 160-400 mg/day with variations depending on food culture and major food sources. Dairy foods remain a food of choice for use as delivery vehicle for many functional ingredients including phytosterols and there are many dairy products available in the global markets which are enriched with phytosterols. The use of phytosterols in commonly consumed dairy products may soon provide an effective tool against CVD and its introduction to our food products is worth anticipating in the near future as nutraceuticals for healthy ageing.

DOI: 10.4018/978-1-5225-3267-5.ch006

INTRODUCTION

Cardiovascular diseases were collectively the number one cause of death in the United States in 2007, accounting for one third of deaths (Xu, Kochanek, Murphy, & Tejada-Vera, 2010). Most of these deaths were attributable to ischemic heart diseases, in which a portion of the muscle in the heart does not receive adequate blood flow. A major cause of ischemia is atherosclerosis, which is characterized by inflammation and lipid accumulation in the lumen of arteries, which decreases blood flow by narrowing the artery and can eventually lead to thrombosis, thereby completely blocking the artery with a blood clot. The Framingham Heart Study is often credited with being the first to systematically investigate the links between blood cholesterol concentrations as a risk factor for cardiovascular diseases, and particularly the effects of low-density lipoprotein cholesterol (LDL-C), highdensity lipoprotein cholesterol (HDL-C), and total cholesterol (Castelli, 1988). The Framingham Heart Study, a long-term observational study, is now observing the third generation of participants from residents in Framingham, MA. The data from this study have allowed a number of prediction models of cardiovascular disease risk associated with a variety of risk factors, and has worked to separate environmental, genetic, and other effects through epidemiological modeling. In one such model, cholesterol concentrations were determined to be such a significant risk factor for coronary heart disease that all other risk factors, such as diabetes, smoking, and blood pressure, were converted into either cholesterol or LDL-C risk points, which were then compared to categories on a risk chart (Wilson, D'Agostino, Levy, Belanger, Silbershatz, & Kannel, 1998). Clearly, LDL-C concentrations are an important risk factor for heart disease.

Cholesterol in the body mainly comes from two sources- endogenous synthesis and dietary consumption. Dietary cholesterol contributes a relatively minor proportion of the cholesterol in the system, with an average of only 276 mg of cholesterol consumed per day (Agricultural Research Service USDA, 2010). In addition, dietary cholesterol minimally affects blood cholesterol, with estimates of only a 0.022-0.027 mg/dL difference in blood cholesterol for every 1 mg of dietary cholesterol consumed (McNamara, 2000). To put this into perspective, an individual with high cholesterol (>240 mg/dL) would need to consume an additional 100 mg of cholesterol per 6 day to increase their blood cholesterol by about 1%. Although dietary cholesterol alone seems to have little effect on blood cholesterol, dietary cholesterol is not the only source of cholesterol per day into the small intestine. Of the total cholesterol in the intestinal lumen, approximately 50% is absorbed, making inhibition of intestinal cholesterol absorption, not just decreasing dietary cholesterol, an important target for lowering blood cholesterol concentrations (McNamara, 2000).

23 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-

global.com/chapter/nutraceutical-intervention-of-phytosterols-

in-cardiovascular-aging/207973

Related Content

Wireless Heartrate Monitoring Along Prioritized Alert Notification Using Mobile Techniques

Rajasekaran Rajkumar (2019). International Journal of Applied Research on Public Health Management (pp. 35-46).

www.irma-international.org/article/wireless-heartrate-monitoring-along-prioritized-alertnotification-using-mobile-techniques/218867

Telemedicine's Role in Pandemic Response and Control Measures

Sabakun Naher Shetuand Takrima Jannat (2022). International Journal of Applied Research on Public Health Management (pp. 1-18). www.irma-international.org/article/telemedicines-role-in-pandemic-response-and-controlmeasures/309410

E-Commerce and IT Projects: Evaluation and Management Issues in Australian and Taiwanese Hospitals

Wenqi Jacintha Hee, Geoffrey Jalleh, Hung-Chih Laiand Chad Lin (2017). International Journal of Public Health Management and Ethics (pp. 69-90). www.irma-international.org/article/e-commerce-and-it-projects/193583

Magnetocaloric as Solid-State Cooling Technique for Energy Saving

Ciro Aprea, Adriana Greco, Angelo Maiorinoand Claudia Masselli (2020). *Global Issues and Innovative Solutions in Healthcare, Culture, and the Environment (pp. 226-252).*

www.irma-international.org/chapter/magnetocaloric-as-solid-state-cooling-technique-for-energysaving/254995

IoT and Big Data in Public Health: A Case Study in Colombia

Fernando Enrique Lopez Martinez, Maria Claudia Bonfante, Ingrid Gonzalez Artetaand Ruby Elena Muñoz Baldiris (2021). *Research Anthology on Public Health Services, Policies, and Education (pp. 399-409).*

www.irma-international.org/chapter/iot-and-big-data-in-public-health/281985