Chapter 7 Dietary Fibers and Nutraceuticals in Prevention of Hypertension

John Intru Disouza

Tatyasaheb Kore College of Pharmacy, India

Kiran Shivaji Patil

Tatyasaheb Kore College of Pharmacy, India

Pratik Shailendra Kakade

Institute of Chemical Technology, India

Vandana Bharat Patravale

Institute of Chemical Technology, India

ABSTRACT

Hypertension is the major cause of mortality amongst many cardiovascular risk factors causing 7.5 million deaths annually. Macronutrient and micronutrient deficiencies are very common in general population and have broad-ranging physiological effects in-vivo which lessen inflammatory cascades and vascular reactivity. A recent trend is to perform nutritional epidemiological studies linking overall diet pattern to the lifestyle, examining the link between food and nutrients of diet to risk of chronic diseases. This chapter would deal with pharmacological and pathological basis of hypertension, utilization of dietary fibers, functional foods, nutraceuticals for hypertensive populations as well as to those with increased cardiovascular risks.

DOI: 10.4018/978-1-5225-3267-5.ch007

INTRODUCTION

Cardiovascular diseases (CVD) are the leading cause of death globally accounting for more than 20 million deaths annually. 3.9 million Deaths in Europe and over 1.8 million deaths in the European Union (EU) occur due to (CVD) every year (European Cardiovascular Disease Statistics, 2017; EHN - European Heart Network, n.d.). Cardiovascular diseases are a group of disorders of the heart and blood vessels. People who are at high cardiovascular risk due to the presence of risk factors such as hypertension, diabetes, hyperlipidemia or already established disease, need early detection and management using counseling and medicines, as appropriate.

HYPERTENSION

Hypertension is a major risk factor for morbidity and mortality associated with stroke, heart failure and coronary artery disease and chronic kidney disease (CKD) (Mancia, Fagard, Narkiewicz, Redón, Zanchetti, & Böhm, 2013). Hypertension in adults of age 18 years and older is defined as systolic blood pressure (SBP) of 140 mmHg or greater and/or diastolic blood pressure (DBP) of 90 mmHg or greater or any level of blood pressure in patients taking antihypertensive medication (Chobanian, Bakris, Black, Cushman, Green, & Izzo, 2003).

CVD nevertheless remains the main cause of death, especially ischemic heart disease and cerebrovascular disease (CD)(Kearney et al., 2005; Zimmet, Alberti, & Shaw, 2001; Jackson, Lawes, Bennett, Milne, & Rodgers, 2005). Hypertension, was identified in a recent World Health Organization report as among the most important preventable causes of premature death. Atherosclerosis is the basic reason for high blood pressure, in which plaque builds inside the arteries. Several factors which lead to atherosclerosis are faulty food habits, lack of exercise, stress, sedentary lifestyle, etc. (McGill, 1979; He, Merchant, Rimm Rosner, Stampfer, Willett, & Ascherio,

Table 1. Types of hypertension

Category Pressure	Systolic Pressure mm Hg	Diastolic Pressure mm Hg
• Normal	90–119	60-79
Pre-hypertension	120–139	80–89
• Stage 1	140–159	90–99
• Stage 2	≥160	≥100
Isolated systolic Hypertension	≥140	<90

39 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-publisher

global.com/chapter/dietary-fibers-and-nutraceuticals-inprevention-of-hypertension/207975

Related Content

Social Challenges Faced by Health Service Providers in Managing COVID-19

S. S. M. Sadrul Huda, Segufta Dilshad, Hamida Mosharrafand Md. Ishtiak Uddin (2022). *International Journal of Applied Research on Public Health Management (pp. 1-6).*

 $\underline{\text{www.irma-international.org/article/social-challenges-faced-by-health-service-providers-inmanaging-covid-19/313439}$

A Review on Chemsensors: A Versatile Tool to Detect Environmentally Toxic Metal Ions

Richa Saxenaand Shweta Saxena (2019). *Environmental Exposures and Human Health Challenges (pp. 133-146).*

www.irma-international.org/chapter/a-review-on-chemsensors/225870

Automation of Intraoperative Analysis of the Indicators of the Inflammatory Response of Neurosurgical Patients Undergoing Brain Tumor Removal: Information Processing, Analysis, Prognosis

Sergey Sokolov, Anatoliy Kondratiev, Nataliya Lestevaand Nataliya Dragina (2023). *International Journal of Applied Research on Public Health Management (pp. 1-13).* www.irma-international.org/article/automation-of-intraoperative-analysis-of-the-indicators-of-the-inflammatory-response-of-neurosurgical-patients-undergoing-brain-tumor-removal/325652

Design of Nano-Scale Devices Affecting Synapses: The New Approach to Artificial Intelligence and Brain Interface

Rinat Galiautdinov (2019). *International Journal of Applied Nanotechnology Research* (pp. 66-78).

www.irma-international.org/article/design-of-nano-scale-devices-affecting-synapses/258911

Gender-Based Differing Effects of Smoking Addiction on COVID-19 Performance

Adem Gökand Nausheen Sodhi (2022). Contemporary Issues in Global Medicine and Moving Toward International Healthcare Equity (pp. 306-316).

 $\underline{www.irma\text{-}international.org/chapter/gender-based-differing-effects-of-smoking-addiction-on-covid-19-performance/312284}$