

Chapter 1

A System to Measure Physiological Response During Social Interaction in VR for Children With ASD

Karla Conn Welch

University of Louisville, USA

Zachary E. Warren

Vanderbilt University, USA

Uttama Lahiri

*Indian Institute of Technology
Gandhinagar, India*

Nilanjan Sarkar

Vanderbilt University, USA

ABSTRACT

This chapter presents work aimed at investigating interactions between virtual reality (VR) and children with autism spectrum disorder (ASD) using physiological sensing of affective cues. The research objectives are two-fold: 1) develop VR-based social communication tasks and integrate them into the physiological signal acquisition module to enable the capture of one's physiological responses in a time-synchronized manner during participation in the task and 2) conduct a pilot usability study to evaluate a VR-based social interaction system that induces an affective response in ASD and typically developing (TD) individuals by using a physiology-based approach. Physiological results suggest there is a different physiological response in the body in relation to the reported level of the affective states. The preliminary results from a matched pair of participants could provide valuable information about specific affect-eliciting aspects of social communication, and this feedback could drive individualized interventions that scaffold skills and improve social wellbeing.

DOI: 10.4018/978-1-5225-7467-5.ch001

INTRODUCTION

Autism is a neurodevelopmental disorder characterized by core deficits in social interaction and communication accompanied by restricted patterns of interest and behavior (American Psychiatric Association, 2000). Currently, with Centers for Disease Control and Prevention (CDC) prevalence estimates for the broad autism spectrum as high as 1 in 59 children (Baio et al., 2018), identification and effective treatment of autism spectrum disorder (ASD) is often characterized as a public health emergency (Interagency Autism Coordinating Committee, 2009). While there is at present no single accepted intervention, treatment, or known cure for ASD, there is growing consensus that intensive behavioral and educational intervention programs can significantly improve short and long term outcomes for individuals with ASD and their families (Cohen, Amerine-Dickens, & Smith, 2006; National Research Council, 2001; Rogers, 1998). While such intervention paradigms have demonstrated significant effect in addressing basic early deficits in young children (i.e., preschool children), traditional interventions designed to address higher level social and adaptive impairments (i.e., communicating with others, processing and integrating information from the environment, establishing and sustaining social relationships, participating in new environments, learning skills related to functional independence) have been demonstrated to be minimally effective for school-aged children and adolescents with ASD. Specifically, low treatment effects and low generalization effects are typical for social and adaptive skill interventions and are thought to be a result of a failure of traditional methodologies to systematically match intervention strategies to specific skill deficits within and across naturalistic settings in appropriately intensive dosages (Bellini, Peters, Benner, & Hopf, 2007). With a lack of widely available efficacious treatment modes for addressing these complex skills at later points in childhood, adolescence, and early adulthood, it is not surprising that evidence suggests a majority of individuals with ASD (i.e., >70%) fail to achieve adaptive independence as adults (Billstedt, Gillberg, & Gillberg, 2005; Cederlund et al., 2007). Current intervention limitations are compounded by the lack of available resources in many areas and the fact that many children are not identified with ASD until later ages (Christensen et al., 2016; Croen, Grether, & Selvin, 2002; Mandell, Listerud, Levy, & Pinto-Martin, 2002; Yeargin-Allsopp et al., 2003). Thus, despite the urgent need and societal import of intensive treatment of social and adaptive impairment at later ages, effective and appropriate intervention resources for children,

31 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/chapter/a-system-to-measure-physiological-response-during-social-interaction-in-vr-for-children-with-asd/227270

Related Content

Web Accessibility: Current Trends

Simon Harper and Yeliz Yesilada (2011). *Handbook of Research on Personal Autonomy Technologies and Disability Informatics* (pp. 172-190).

www.irma-international.org/chapter/web-accessibility-current-trends/48281

Clustering Methods for Gene-Expression Data

L.K. Flack (2009). *Handbook of Research on Systems Biology Applications in Medicine* (pp. 209-220).

www.irma-international.org/chapter/clustering-methods-gene-expression-data/21533

Role of Acoustic Properties in Biomedical Active Noise Control

Sajil C. K. and Achuthsankar S. Nair (2020). *International Journal of Biomedical and Clinical Engineering* (pp. 48-60).

www.irma-international.org/article/role-of-acoustic-properties-in-biomedical-active-noise-control/240746

Methods and Applications for Segmenting 3D Medical Image Data

Hong Shen (2009). *Medical Informatics: Concepts, Methodologies, Tools, and Applications* (pp. 1144-1161).

www.irma-international.org/chapter/methods-applications-segmenting-medical-image/26287

A WBAN-Based Framework for Health Condition Monitoring and Faulty Sensor Node Detection Applying ANN

Koushik Karmakar, Sohail Saif, Suparna Biswas and Sarmistha Neogy (2021). *International Journal of Biomedical and Clinical Engineering* (pp. 44-65).

www.irma-international.org/article/a-wban-based-framework-for-health-condition-monitoring-and-faulty-sensor-node-detection-applying-ann/282494