Chapter 39 Bioremediation: New Prospects for Environmental Cleaning by Enzymes

Vikram Singh

Agricultural Research Organization, Volcani Center, Israel

ABSTRACT

Pollution is the biggest menace to the living being in this planet today. Enzyme bioremediation is a "breakthrough technology" that holds the potential of pollutant eradication through exploiting the enzyme potential by using the various techniques. Enzyme biocatalysis is referred as white biotechnology and work by green chemistry concept. Moreover, developments in the design and application of enzyme cocktails, mutienzyme complexes, promiscuous enzymes and protein families (cupin and VOC superfamily) has recently emerged a new opportunity in bioremediation. The implementation of various enzyme modification approaches intended for potential bioremediation has been done by adopting enzyme immobilization using magnetic nanoparticles, designer enzymes generation through enzyme engineering, nano-technological advancement for single enzyme nanoparticle generations, electro-bioremediation and carbon nanotube construction. Hence, enzyme bioremediation have greater positive effects and propose significant promise to pollutant bioremediation. In conclusion, the enzymatic bioremediation open the new era of pollutant eradication for clean, safe and green environment.

INTRODUCTION

Environmental pollution is one of the serious issues of the present and future climatic scenario. The soaring environmental pollution affects the all forms of living and non-living globally. Environmental contamination has become a serious problem with the advancement of industrial revolution in developed and producing nations. The increasing population growth and urbanization stretch the use of innate resources for the maximum yield. The carrying capacity of the earth is significantly smaller than the demands with increasing populations. To satisfy the existing demands, the overuse of natural resources often results in nature's degradation (Santos, 1990). The new agricultural practices have been carried out

DOI: 10.4018/978-1-5225-8903-7.ch039

Bioremediation

to meet the food demand of soaring population. Pesticides play a significant part in modern agriculture to increase the productivity. However, pesticides may have a good effect on agricultural productivity, conversely, their indiscriminate use in these farming practices has contributed to serious concern for the health of living being and the environment. (Diez, 2010; Coutinho et al., 2005). Pesticide poisonings are a massive problem worldwide, especially in developing countries. Out of the million tons of pesticides applied annually, most of them get accumulated in the environment with the degradation of only a small fraction (Eerd, Hoagland, Zablotowicz, & Hall, 2003). Pesticides are recalcitrant and remain for a long time in soils and enter directly into the food chain. To overcome these problems researcher has released the new degradable pesticides. The fate of pollutant in the environment is influenced by biological, chemical and physical degradation processes which determine their persistence and mobility (Gavrilescu, 2005). Consideration of the current technology for removal of pollutants clearly shows that physical and chemical methods are often uneconomical. Nevertheless, biological treatment technology may provide a resolution to this problem. The pollutant degradation by biological means such as microorganisms (Microbioremediation) or plant (Phytoremediation) etc. is termed as "Bioremediation". The use of bioremediation technologies provides a dependable and economic alternative to commonly used physical and chemical handling. Biological remediation is regarded as the most dependable, least disruptive and most cost-effective treatment method (Mueller, Cerniglia, & Pritchard, 1996). Principle of bioremediation is to utilize the pollutants, use them for their growth and metabolism or convert them from toxic to nontoxic form. Bioremediation by microorganism offer more advantage over phytoremediation as microbioremediation is a well established technology for the pollutant removal. Simultaneously, microbes act as an 'eco-friendly nano-factories' for environmental cleaning. Microbioremediation offers natural, environmentally friendly and economical solution for environmental cleaning (Vidali, 2001). Bacteria and fungi are well known to degrade hazardous organic contaminants to environmentally less toxic or nontoxic compounds. These organic compounds are readily incorporated into the microbes and get oxidized under aerobic or anaerobic conditions. Heterotrophic bacteria and fungi are the chief agents causing biodegradation. Recently, genetically modified microorganisms have been directed to formulate bioremediation (Paul, Pandey, Pandey, & Jain, 2005). The Pseudomonas fluorescens HK44 possesses a naphthalene catabolic plasmid is the first genetically engineered microorganism approved for bioremediation testing (Ford, Savler, & Burlage, 1999). The major problems associated with microbioremediation is the mass transfer, additionally they also require aeration and nutrient on contaminated sites (require bio-stimulation) and also suffer from thermal condition problems (Fantroussi & Agathos, 2005; Scow & Hicks, 2005). Now a day the bioremediation is shifted from microbes to enzymes because of their ecofriendly attributes. The pollutant degradation by an enzyme is known as "enzymatic bioremediation" and particularly suited for rapid remediation. This method is adventitious over to microbial bioremediation (either natural or genetically engineered microbes), because of the efficiency, time consumption and expense (Ahuja, Ferreira, & Moreira, 2004). The exploited enzyme potential using the various techniques of in-vitro enzyme evolution is being revolutionized in present scenario. The enzymatic bio-efficiency can be appraised by the adoption of two recent methods Biostimulation and Bioaugmentation (Sutherland, Russell, & Selleck, 2002; Scott et al., 2008). Biostimulation provides the optimal environment to enzymes or microbes for their maximum efficiency" in terms of temperature, pH, moisture, aeration, etc. (Baker & Herson, 1994), however "Bioaugmentation employ the superior biodegradable agent (enzymes or microbes) for the environment cleaning" (Scragg, 2005). Enzymes are having narrow (chemo, region and stereo selectivity) or broad specificity and can be applied to a vast range of compounds effectively 27 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/bioremediation/228656

Related Content

Impact of Diesel-Butanol-Waste Cooking Oil Biodiesel Blends on Stationary Diesel Engine Performance and Emission Characteristics

H. Sharon, Joel Jackson R.and Prabha C. (2020). *Recent Technologies for Enhancing Performance and Reducing Emissions in Diesel Engines (pp. 173-192).*

www.irma-international.org/chapter/impact-of-diesel-butanol-waste-cooking-oil-biodiesel-blends-on-stationary-dieselengine-performance-and-emission-characteristics/249063

Applications of Supercomputers in Population Genetics

Gerard G. Dumancas (2019). *Biotechnology: Concepts, Methodologies, Tools, and Applications (pp. 693-719).*

www.irma-international.org/chapter/applications-of-supercomputers-in-population-genetics/228645

Biosynthesis of Silver Nanoparticles for Study of Their Antimicrobial Effect on Plasma-Treated Textiles: Silver Coating of Plasma-Treated Cotton Fabric

Shazia Shukrullah, Muhammad Anwar, Muhammad Yasin Nazand Inzamam UI Haq (2022). *Emerging Developments and Applications of Low Temperature Plasma (pp. 149-166).*

www.irma-international.org/chapter/biosynthesis-of-silver-nanoparticles-for-study-of-their-antimicrobial-effect-on-plasmatreated-textiles/294715

Microbial Degradation of Azo Dyes: The Role of Azoreductase to Initiate Degradation

Dirk Tischler, Jingxian Qi, Anna Christina R. Ngoand Michael Schlömann (2019). *Biotechnology: Concepts, Methodologies, Tools, and Applications (pp. 1867-1897).*

www.irma-international.org/chapter/microbial-degradation-of-azo-dyes/228696

Domain-Based Approaches to Prediction and Analysis of Protein-Protein Interactions

Morihiro Hayashidaand Tatsuya Akutsu (2019). *Biotechnology: Concepts, Methodologies, Tools, and Applications (pp. 406-427).*

www.irma-international.org/chapter/domain-based-approaches-to-prediction-and-analysis-of-protein-protein-interactions/228632