Chapter 7 Cognitive Enhancement and Neuroprotective Abilities of Plants: Ethnobotanical and Pharmacological Importance of These Plants

Kiran Mustafa The Women University Multan, Pakistan

Javaria Kanwal The Women University Multan, Pakistan

Samia Khakwani The Women University Multan, Pakistan

Sara Musaddiq The Women University Multan, Pakistan

ABSTRACT

Extensive research suggests that a number of plant-derived chemicals and traditional Oriental herbal remedies possess cognition-enhancing properties. Widely used current treatments for dementia include extracts of Ginkgo biloba and several alkaloidal, and therefore toxic, plant-derived cholinergic agents. Several non-toxic, European herbal species have pan-cultural traditions as treatments for cognitive deficits, including those associated with aging. Acute administration has also been found to reliably improve mnemonic performance in healthy young and elderly

DOI: 10.4018/978-1-7998-2094-9.ch007

Cognitive Enhancement and Neuroprotective Abilities of Plants

cohorts, whilst a chronic regime has been shown to attenuate cognitive declines in sufferers from Alzheimer's disease. The present chapter looks at the ethnobotanical and pharmacological importance of various plants cognitive enhancing and other neuroprotective abilities.

INTRODUCTION

Gradual loss of function or structure of the nerve cells lead to the neurodegenerative disorders such as Parkinson disease, Alzheimer disease and Huntington's disease, multiple scleroses etc. According to World Health Organization (WHO), the second principal cause of death will be neurodegenerative diseases by the year 2040 (Tejada et al., 2017). Recently immense focused has been placed on understanding the mechanism of death of nerve cells. Neurotic injuries are associated with the series of neural toxic agents including inflammatory cytokines, mitochondrial malfunctioned, oxidative stress and irregular protein activities (Ansari & Khodagholi, 2013; Chen et al., 2016; Hirsch & Hunot, 2009). One of the leading reasons behind neuron death is reactive oxygen species (ROS) that induced oxidative harm to protein, lipids and DNA followed by persuading oxidative strain. Besides this oxidative stress is predominantly associated with death of secondary cells in various central nervous system ailments (Jalsrai et al., 2016; Sultana & Butterfield, 2010). Recent analysis has shown that neuron death is occurred by apoptosis. There are two major signaling pathways that participated in the apoptotic cell death; mitochondrial pathway (intrinsic) and death of receptors (extrinsic) (Elmore, 2007). Different transcription factors are involved in neuron protection, such as cAMP-response element binding protein, nuclear factor erythroid-derived 2, nuclear-factor-kappa-B etc. These transcriptional agents are primarily linked with regulation of various genes that modulate the inflammatory action and activation of antioxidant enzyme (Ansari et al., 2011; Ashabi et al., 2012). Additionally, transcription factors are because of their functions in growth of nerve cells, synaptic plasticity, axonogenesis, and neuronal homeostasis (Freese et al., 2010). In this context, neuroprotective elements appeared as a shielding agent to secure CNS against disorders associated to neuron (Pak et al., 2016). Various synthetic drugs are present for neurodegenerative disorders like acetylcholinesterase blockers, neuroleptic drugs, dopamine therapy, brain stimulation, antipsychotic drugs (Mizuno, 2014) non-steroidal anti-inflammatory drugs and riluzole etc. (Chen & Pan, 2014). These drugs are useful against several neurodegenerative diseases but possess

23 more pages are available in the full version of this document, which may be purchased using the "Add to Cart"

button on the publisher's webpage: www.igi-

global.com/chapter/cognitive-enhancement-and-

neuroprotective-abilities-of-plants/252939

Related Content

Therapeutic Importance and Application of Boswellic Acid From the Plant Boswellia serrata

Raghunath Satpathy (2020). Advanced Pharmacological Uses of Medicinal Plants and Natural Products (pp. 302-315).

www.irma-international.org/chapter/therapeutic-importance-and-application-of-boswellic-acidfrom-the-plant-boswellia-serrata/252951

Scoring Functions in Docking Experiments

Pravin Ambureand Kunal Roy (2016). *Methods and Algorithms for Molecular Docking-Based Drug Design and Discovery (pp. 54-98).* www.irma-international.org/chapter/scoring-functions-in-docking-experiments/151883

Quantitative Nanostructure-Activity Relationship Models for the Risk Assessment of NanoMaterials

Eleni Vrontaki, Thomas Mavromoustakos, Georgia Melagrakiand Antreas Afantitis (2017). *Pharmaceutical Sciences: Breakthroughs in Research and Practice (pp. 1314-1338).*

www.irma-international.org/chapter/quantitative-nanostructure-activity-relationship-models-forthe-risk-assessment-of-nanomaterials/174171

The Comparison of Docking Search Algorithms and Scoring Functions: An Overview and Case Studies

Marjana Novi, Tjaša Tibaut, Marko Anderluh, Jure Borišekand Tihomir Tomaši (2017). *Pharmaceutical Sciences: Breakthroughs in Research and Practice (pp. 820-849).*

www.irma-international.org/chapter/the-comparison-of-docking-search-algorithms-and-scoringfunctions/174152

Mushroom Polysaccharides as Biological Response Modifiers in Cancer Therapy

Meera C.R. (2023). *Natural Products as Cancer Therapeutics (pp. 176-195).* www.irma-international.org/chapter/mushroom-polysaccharides-as-biological-responsemodifiers-in-cancer-therapy/329159