

## Chapter 2

# Higher Dimensions of Clusters of Intermetallic Compounds

### ABSTRACT

*The author has previously proved that diffraction patterns of intermetallic compounds (quasicrystals) have translational symmetry in the space of higher dimension. In this chapter, it is proved that the metallic nanoclusters also have a higher dimension. The internal geometry of clusters was investigated. General expressions for calculating the dimension of clusters are obtained from which it follows that the dimension of metallic nanoclusters increases linearly with increasing number of cluster shells. The dimensions of many experimentally known metallic nanoclusters are determined. It is shown that these clusters, which are usually considered to be three-dimensional, have a higher dimension. The Euler-Poincaré equation was used, and the internal geometry of clusters was investigated.*

### INTRODUCTION

A systematic study of the geometry of the structures of chemical compounds (Zhizhin, 2014a, b, c, d, 2016, 2018, 2019a) showed that almost all elements of the periodic system form molecules of higher dimension. It is natural to assume that clusters, as larger than education molecules, including a large number of atoms, can have a higher dimension. However, until recently, clusters considered as three - dimensional objects (Lord, Mackay, & Ranganathan, 2006; Pauling, 1960). This Chapter discusses clusters of real chemical


DOI: 10.4018/978-1-7998-3784-8.ch002

compounds (as opposed to the book of Diudea, 2018, in which, as the author emphasizes, clusters are not related to real chemical compounds). Moreover, in this work, consideration of clusters is limited to a special type of chemical compounds - intermetallic compounds, since the study of intermetallic compounds has had a significant impact on the development of scientific views in recent decades. In particular, the discovery of so - called quasicrystals is associated with intermetallic compounds, i.e. crystals supposedly devoid of translational symmetry (Shechtman et al., 1984). Although it was later shown that quasicrystals have translational symmetry, but in the space of higher dimension (Shevchenko, Zhizhin, & Mackay, 2013a, b; Zhizhin, 2014c; Zhizhin, & Diudea, 2016). The ideas about the higher dimension of clusters and the calculation of this dimension should be taken account to in their practical use as objects with valuable physic - chemical properties.

## **CLUSTERS OF MACKAY**

Mackay's cluster consists of two icosahedrons of different sizes with a common center (Mackay, 1962). A larger icosahedron is obtained by attaching a number of tetrahedrons and octahedrons to the surface of the smaller icosahedron.

*Figure 1. The cluster of Mackay*



25 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: [www.igi-global.com/chapter/higher-dimensions-of-clusters-of-intermetallic-compounds/261001](http://www.igi-global.com/chapter/higher-dimensions-of-clusters-of-intermetallic-compounds/261001)

## Related Content

---

### A Reduced-Order General Continuum Method for Dynamic Simulations of Carbon Nanotube

Yang Yang and William W. Liou (2010). *International Journal of Nanotechnology and Molecular Computation* (pp. 1-24).

[www.irma-international.org/article/reduced-order-general-continuum-method/52086](http://www.irma-international.org/article/reduced-order-general-continuum-method/52086)

### Simple Collision-Based Chemical Logic Gates with Adaptive Computing

Rita Toth, Christopher Stone, Ben de Lacy Costello, Andrew Adamatzky and Larry Bull (2011). *Theoretical and Technological Advancements in Nanotechnology and Molecular Computation: Interdisciplinary Gains* (pp. 162-175).

[www.irma-international.org/chapter/simple-collision-based-chemical-logic/50141](http://www.irma-international.org/chapter/simple-collision-based-chemical-logic/50141)

### Designing Biomedical Stents for Vascular Therapy: Current Perspectives and Future Promises

Arghya Paul (2014). *Nanotechnology: Concepts, Methodologies, Tools, and Applications* (pp. 972-996).

[www.irma-international.org/chapter/designing-biomedical-stents-for-vascular-therapy/102052](http://www.irma-international.org/chapter/designing-biomedical-stents-for-vascular-therapy/102052)

### A New Approach for DNA Sequence Similarity Analysis based on Triplets of Nucleic Acid Bases

Dan Wei, Qingshan Jiang and Sheng Li (2010). *International Journal of Nanotechnology and Molecular Computation* (pp. 1-11).

[www.irma-international.org/article/new-approach-dna-sequence-similarity/53349](http://www.irma-international.org/article/new-approach-dna-sequence-similarity/53349)

### Cluster Origin of Solvent Features of Fullerenes, Single-Wall Carbon Nanotubes, Nanocones, and Nanohorns

Francisco Torrens and Gloria Castellano (2012). *Nanoscience and Advancing Computational Methods in Chemistry: Research Progress* (pp. 1-57).

[www.irma-international.org/chapter/cluster-origin-solvent-features-fullerenes/66244](http://www.irma-international.org/chapter/cluster-origin-solvent-features-fullerenes/66244)