Chapter 4 Physicochemical Characteristics of Indonesian Native Starch

Maria Belgis University of Jember, Indonesia

Nur Fathonah Sadek Bina Nusantara University, Indonesia

Ardiyan Dwi Masahid University of Jember, Indonesia

Dian Purbasari University of Jember, Indonesia

Dyah Ayu Savitri University of Jember, Indonesia

ABSTRACT

Indonesia has many local starch sources, including rice, taro, arrowroot, sago, corn, konjac, cassava, and elephant foot yam (suweg). Several uses of natural starch have been widely used as the primary raw material or as a food additive. Natural starch generally has some drawbacks limiting its use, such as color, solubility index, retrogradation, and paste stability. Starch modification is needed to improve these limitations. Starch modification is carried out physically and chemically. In this chapter, the authors will discuss the physical and chemical characteristics of several local Indonesian starches, both natural and modified, and their potential to be developed as food ingredients.

DOI: 10.4018/978-1-6684-5629-3.ch004

This chapter published as an Open Access Chapter distributed under the terms of the Creative Commons Attribution License (http:// creativecommons.org/licenses/by/4.0/) which permits unrestricted use, distribution, and production in any medium, provided the author of the original work and original publication source are properly credited.

INTRODUCTION

Starch is a carbohydrate polymer used as the primary source of calories for consumption. Apart from being used as a food ingredient, starch may act as a gelling agent, thickener, foam and emulsion stabilizer, battered and breaded, binder, crispy texture maker, clouding stabilizer, and flavor encapsulant. Starch could also be employed in the paper, textile, and pharmaceutical industries. Various plant sources of starch, such as rice, taro, arrowroot, sago, corn, konjac, cassava, and elephant foot yam (suweg), are growing a lot in Indonesia. They respectively produce starch with different properties. This chapter reviews the physicochemical characterization of several Indonesian starch plants and their roles in controlling food processing. The scientific information from this chapter is expected to enrich the study of starch sources worldwide.

Rice (Oryza sativa)

Rice is a carbohydrate source and a staple food for most Indonesians. Rice is mainly consumed as cooked polished grains, whereas rice flour is used as an ingredient in many Indonesian cuisines. Starch is the most significant component in rice grains and determines the quality of rice products. The starch content in rice is in the range of 80% (Hasjim et al., 2013).

Gelatinization characteristic is considered one of the essential properties in evaluating rice starch properties. In particular, gelatinization is the most crucial rheological indicator of the quality characteristics of rice cooking and processing. Amylose content and branch chain length distribution have also been found to impact rice starch's swelling and pasting properties substantially. Rice starch granules with high amylose content act as swelling inhibitors. Amylopectin contributes to swelling, while amylose and amylose-lipid complex inhibit swelling. In addition, low swelling levels and solubility tend to increase with the increasing temperature of amylose molecules in rice. Studies show that a higher level of long chain amylopectin in rice starch is associated with firmer rice after cooking. Retrogradation was also found higher in rice starch with a longer amylopectin chain (Yoon et al., 2012).

Taro (Colocasia esculenta)

Taro has a starch content of about 80%. The characteristics of taro starch have been evaluated based on swelling power and solubility index. The swelling behavior of starch can be influenced by amylose and amylopectin compositions. According to Aboubakar et al. (2008), the swelling power of taro starch will increase along with starch cooking level, indicated by an increase in Water Absorption Capacity (WAC).

16 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: <u>www.igi-</u> <u>global.com/chapter/physicochemical-characteristics-of-</u> <u>indonesian-native-starch/319451</u>

Related Content

Estimating the Cost of Drained Lands by Using Them in Variable Conditions

Nadiia Frolenkova, Ruslan Tykhenko, Larysa Semenkoand Olha Tykhenko (2023). Handbook of Research on Improving the Natural and Ecological Conditions of the Polesie Zone (pp. 359-371).

www.irma-international.org/chapter/estimating-the-cost-of-drained-lands-by-using-them-invariable-conditions/324049

Enhanced F-Perceptory Approach for Dealing With Geographic Data Imprecision From the Conceptual Modeling to the Fuzzy Geographical Database Building

Besma Khalfi, Cyril De Runzand Herman Akdag (2019). *Environmental Information Systems: Concepts, Methodologies, Tools, and Applications (pp. 426-455).* <u>www.irma-international.org/chapter/enhanced-f-perceptory-approach-for-dealing-with-</u> <u>geographic-data-imprecision-from-the-conceptual-modeling-to-the-fuzzy-geographical-database-</u> <u>building/212953</u>

Future of Public Sector Enterprises in the Metaverse

Richmond Anane-Simonand Sulaiman Olusegun Atiku (2023). *Multidisciplinary Approaches in AI, Creativity, Innovation, and Green Collaboration (pp. 167-188).* www.irma-international.org/chapter/future-of-public-sector-enterprises-in-the-metaverse/322876

Towards a More Gender-Inclusive Climate Change Policy

Farah Kabir (2018). *Climate Change and Environmental Concerns: Breakthroughs in Research and Practice (pp. 525-540).*

www.irma-international.org/chapter/towards-a-more-gender-inclusive-climate-change-policy/201721

Case Study: Evaluation of Climate Change Adaptation Interventions

(2018). Innovative Strategies and Frameworks in Climate Change Adaptation: Emerging Research and Opportunities (pp. 59-82). www.irma-international.org/chapter/case-study/191158