

Chapter XV

Evidence on the Efficacy of Integrated Care

Torben Larsen

University of Southern Denmark, Denmark

ABSTRACT

The fragmented delivery of healthcare and social services was put on the agenda as a major problem by WHO in 2002. Early home-supported discharge (EHSD) of stroke patients combining efficacy with net savings represents a prototype of integrated care (IC) or overlapping services for better clinical continuity. Other frequent chronic conditions as heart failure, chronic obstructive pulmonary disease, and mental disease exhibit parallel results from home health interventions. A SWOT analysis of IC emphasizes: Strengths are 1) economic dominance, 2) benefits to a majority of the population, and 3) psychological motivators for the patient (Hawthorne effect). Weaknesses are 1) moderate improvements on a day-to-day basis, 2) some lack of RCT, and 3) lack of trust across settings. Opportunities are 1) job enrichment to health professionals, 2) low-tech improvements affordable to low and middle income countries, and 3) organisational quality. Threats are 1) fragmented financial conditions, 2) defensive specialists, and 3) mediocre implementations. A meso-strategy of implementation is recommended to EU (FP7): 1) Make a synthesis of existing and ongoing research as a health technology assessment (HTA) of IC in EU for improved interdisciplinary cooperation across the hospital and primary care interface for selected CC. 2) The organisational dimension should focus on the formation of country specific multidisciplinary networks on IC.

INTRODUCTION

A number of studies from the 1990s have focused on healthcare problems related to lack of clinical continuity. The fragmented delivery of health-

care and social services is put on the agenda as a major problem by WHO (Gröne, 2002) and was followed-up by the European Commission and Council (Joint Report, 2003, p.15). However, a FP5 project on integrated health and social care

for older persons (Leischenring, 2004) concludes that ‘Integrated care (IC) ‘by law’ as a top-down implementation will certainly not suffice, and market mechanisms as bottom-up approaches are less likely to improve joint working and the development of shared visions’. What could then be done to overcome the problems of a fragmented delivery of healthcare?

Fragmented delivery and lack of clinical continuity is more relevant to chronic conditions (CC) than to time limited acute episodic care (Holman, 2004).

This chapter aims to review the present state of research on integrated care (IC) specific to chronic conditions (CC) focusing the efficacy regarding Activities of Daily Living (ADL) in order to develop an effective and economic strategy for clinical continuity.

METHODS AND MATERIALS

General Strategy of Evaluation

The essence of IC is ‘overlapping’ services in the secondary/primary interface after discharge in contrast to coordination at the management level alone (Gröne, 2002, p. 2). IC should apply only as far as quality outcomes are improved with the overall aim to improve equitably distributed population health (Gröne, 2002, p. 3). He illustrates the causal relationships derived from IC as reproduced in Figure 1.

Operational quality outcomes are crucial for the design of IC. Mortality is seldom a major indicator for IC. Typically, studies of effectiveness on IC address activities of daily living (ADL) as:

1. Referrals to permanent institutional care (i.e. nursing homes).
2. Independence in ADL as indicated by functional indices as Barthel Index (BI) or Functional Independence Measure (FIM).
3. Shortened length-of-stay at hospitals / less readmissions.

Evaluation of the relationship between IC and outcomes will follow best international practice as formalised in the international operation of Health Technology Assessment. According to the definition of HTA by EUnetHTA:

Health technology assessment (HTA) is a multi-disciplinary process that summarises information about the medical, social, economic and ethical issues related to the use of a health technology in a systematic, transparent, unbiased, robust manner. It aims to inform the formulation of safe and effective health policies that are patient focused and seek best value.

HTA may address direct and intended consequences of technologies as well as indirect and unintended consequences. The main purpose of HTA is to assist informed technology-based policymaking in health care. Most health professionals and many decision-makers in health care might comply with the conclusions from an HTA as far as it investigates all of the following aspects:

1. Effectiveness regarding the physiologic outcome of the intervention
2. Patient safety and satisfaction, solicited i.e. by focus group interviews

Figure 1. Action model for integrated care (IC)

Population ?	Quality ?	Continuity of care ?	Integrated care
	-access	(as experienced by	(vision, model
	-satisfaction	staff and patients)	strategy)
	-efficiency		
	-effectiveness		

Source: Trends in Integrated Care–Reflections on Concept. Issues, WHO (Gröne, 2002).

15 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/evidence-efficacy-integrated-care/35780

Related Content

Towards Semantic Interoperability in Health Data Management Facilitating Process Mining

Barbara Traxler, Emmanuel Helm, Oliver Krauss, Andreas Schuler and Josef Kueng (2018). *International Journal of Privacy and Health Information Management* (pp. 1-12).

www.irma-international.org/article/towards-semantic-interoperability-in-health-data-management-facilitating-process-mining/211973

Methodology to Set Regulations for Safe Reuse of Wastewater and Sludge for Agriculture in Developing Countries Based on a Scientific Approach and Following the New WHO Guidelines

B. Jimenez and I. Navarro (2009). *Handbook of Research on Information Technology Management and Clinical Data Administration in Healthcare* (pp. 690-709).

www.irma-international.org/chapter/methodology-set-regulations-safe-reuse/35808

Risk-Benefit Analysis of Combination Versus Unopposed HRT in Post-Menopausal Women

Geer Mohammad Ishaq, Pz Tasaduq Hussain, Mir Javid Iqbal and Mohsin Bin Mushtaq (2011). *International Journal of User-Driven Healthcare* (pp. 61-76).

www.irma-international.org/article/risk-benefit-analysis-combination-versus/61323

Key Health Information Systems Outsourcing Issues from Six Hospital Cases

Chad Lin, Yu-An Huang, Chien-Fa Li and Geoffrey Jalleh (2013). *User-Driven Healthcare: Concepts, Methodologies, Tools, and Applications* (pp. 824-837).

www.irma-international.org/chapter/key-health-information-systems-outsourcing/73867

A Two-Stage Registration Method for Dental Volumetric Data and Mesh Data

Shuai Yang, Wenjing Shi, Yongzhen Ke and Yongjiang Xue (2021). *International Journal of Healthcare Information Systems and Informatics* (pp. 1-15).

www.irma-international.org/article/a-two-stage-registration-method-for-dental-volumetric-data-and-mesh-data/279330