

Chapter 28

Exploratory Analysis of Fossil– Fuel CO₂ Emissions Time Series Using Independent Component Analysis

Sargam Parmar

Ganpat University, India

Bhuvan Unhelkar

University of Western Sydney & MethodScience, Australia

ABSTRACT

Carbon dioxide (CO₂) is one of the most important gases in the atmosphere, and is necessary for sustaining life on Earth. However, it is also a major greenhouse gas out of the six that contribute to global warming and climate change. During the last decade technologists, economists and sociologists are taking substantial interest in studying the impact of greenhouse phenomenon. Scientists are trying to find solutions to reduce CO₂ emissions by changes in structure of energy production and consumption. Every attempt is being made to use new models and methods to estimate measure and monitor greenhouse gases in the future. Independent Component Analysis (ICA) is a method for automatically identifying a set of underlying factors in a given data set. This chapter describes the use of the ICA algorithm in Environmentally Intelligent (EI) applications. EI applications have a wide ranging responsibilities including collection, analysis and reporting of environmental data related to the organization. ICA algorithm opens up the opportunity to improve the quality of data being analyzed by these EI applications. ICA finds application in several fields of interest and it is a tempting alternative to try ICA on multivariate time series such as a CO₂ emission from fossil fuel for the period 1950 to 2006. This chapter describes the linear mapping of the observed multivariate time series into a new space of statistically independent components (ICs) that might reveal driving mechanisms for CO₂ emissions that may otherwise remain hidden.

DOI: 10.4018/978-1-61692-834-6.ch028

INTRODUCTION

Carbon dioxide (CO₂) is one of the most important gases in the atmosphere, and is necessary for sustaining life on Earth. However, it is also a major greenhouse gas out of the six (<http://www.epa.gov/climatechange/emissions/>) that contribute to global warming and climate change. During the last decade technologists, economists and sociologists are taking substantial interest in studying the impact of greenhouse phenomenon. Scientists are trying to find solutions to reduce CO₂ emissions by changes in structure of energy production and consumption. Every attempt is being made to use new models and methods to estimate measure and monitor greenhouse gases in the future. Independent Component Analysis (ICA) is a method for automatically identifying a set of underlying factors in a given data set. This chapter describes the use of the ICA in Environmentally Intelligent (EI) applications (see Unhelkar and Trivedi, 2009) that will improve the quality of data being analyzed by these EI applications. This rapidly evolving technique is currently finding applications in several fields of interest and it is a tempting alternative to try ICA on multivariate time series such as a CO₂ emission from fossil fuel for the period 1950 to 2006. The key idea here is to linearly map the observed multivariate time series into a new space of statistically independent components (ICs) that might reveal some driving mechanisms that may otherwise remain hidden.

Estimates of the amount of carbon emitted in to the atmosphere from fossil-fuel burning produce can be considered as a basic time-series in this context. Different kinds of time-series have been recorded and studied. Nowadays, all transactions on carbon emitted to the atmosphere from fossil-fuel burning are recorded, leading to a huge amount of data available, either freely or commercially on the Internet.

Furthermore, the stochastic uncertainties inherent in fossil-fuel CO₂ emissions time-series and the theory needed to deal with them make the subject especially interesting not only to economists, but also to statisticians and physicists. Fossil fuel CO₂ emissions systems is a complex evolved dynamic system with high volatility and noise. Due to its irregularity, fossil fuel CO₂ emissions time series forecasting is regarded as a rather challenging task. CO₂ emission estimate systems require more advanced signal processing methods, and correct reception of CO₂ emission time series is more difficult because of several phenomena such as annual global CO₂ emissions from solid fuels, liquid fuels, natural gas, gas flaring, and cement manufacturing.

This chapter describes the use of the Independent Component Analysis (ICA) in Environmentally Intelligent (EI) applications that will improve the quality of data being analyzed by these EI applications. ICA is a method for automatically identifying a set of underlying factors in a given data set. This rapidly evolving technique is currently finding applications in several fields of interest and it is a tempting alternative to try ICA on multivariate time series such as a CO₂ emission from fossil fuel for the period 1950 to 2006. The key idea here is to linearly map the observed multivariate time series into a new space of statistically independent components (ICs) that might reveal some driving mechanisms that otherwise remain hidden.

The chapter is organized as follows. Section 2 provides a background to ICA and a guide on how to estimate the independent components. Section 3 discusses some issues concerning the general application of ICA to CO₂ emission from fossil fuel. Results are shown for global CO₂ emission in general and specifically for India's CO₂ emission in section 4. Finally, section 5 concludes the chapter with a discussion and thoughts on future directions.

17 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/exploratory-analysis-fossil-fuel-co2/48443

Related Content

An Australian Rules Football Club Approach To Green ICT

Jeffrey Phuah (2011). *Handbook of Research on Green ICT: Technology, Business and Social Perspectives* (pp. 348-354).

www.irma-international.org/chapter/australian-rules-football-club-approach/48439

Scenarios and Modeling of Land Use and Cover Changes in Portugal from 1980 to 2040

Sara Santos, Pedro Cabral and Alexander Zamyatin (2015). *International Journal of Agricultural and Environmental Information Systems* (pp. 1-15).

www.irma-international.org/article/scenarios-and-modeling-of-land-use-and-cover-changes-in-portugal-from-1980-to-2040/137160

Information Systems for a Green Organisation

Yogesh Deshpande and Bhuvan Unhelkar (2011). *Handbook of Research on Green ICT: Technology, Business and Social Perspectives* (pp. 116-130).

www.irma-international.org/chapter/information-systems-green-organisation/48423

Towards The Use of Probabilistic Spatial Relation Databases in Business Process Modeling

Haizhou Li, François Pinet and Farouk Toumani (2015). *International Journal of Agricultural and Environmental Information Systems* (pp. 50-62).

www.irma-international.org/article/towards-the-use-of-probabilistic-spatial-relation-databases-in-business-process-modeling/128850

Identification of Associations between Clinical Signs and Hosts to Monitor the Web for Detection of Animal Disease Outbreaks

Elena Arsevska, Mathieu Roche, Pascal Hendriks, David Chavernac, Sylvain Falala, Renaud Lancelot and Barbara Dufour (2016). *International Journal of Agricultural and Environmental Information Systems* (pp. 1-20).

www.irma-international.org/article/identification-of-associations-between-clinical-signs-and-hosts-to-monitor-the-web-for-detection-of-animal-disease-outbreaks/163316