

Chapter 19

Linking Interactome to Disease: A Network–Based Analysis of Metastatic Relapse in Breast Cancer

Maxime Garcia

Inserm, Paoli Calmettes Institute, France

Olivier Stahl

Inserm, Paoli Calmettes Institute, France

Pascal Finetti

Inserm, Paoli Calmettes Institute, France

Daniel Birnbaum

Inserm, Paoli Calmettes Institute, France

François Bertucci

Inserm, Paoli Calmettes Institute, France

Ghislain Bidaut

Inserm, Paoli Calmettes Institute, France

ABSTRACT

The introduction of high-throughput gene expression profiling technologies (DNA microarrays) in molecular biology and their expected applications to the clinic have allowed the design of predictive signatures linked to a particular clinical condition or patient outcome in a given clinical setting. However, it has been shown that such signatures are prone to several problems: (i) they are heavily unstable and linked to the set of patients chosen for training; (ii) data topology is problematic with regard to the data dimensionality (too many variables for too few samples); (iii) diseases such as cancer are provoked by subtle misregulations which cannot be readily detected by current analysis methods. To find a predictive signature generalizable for multiple datasets, a strategy of superimposition of a large scale of protein-protein interaction data (human interactome) was devised over several gene expression datasets (a total of 2,464 breast cancer tumors were integrated), to find discriminative regions in the interactome (subnetworks) predicting metastatic relapse in breast cancer. This method, Interactome-Transcriptome

DOI: 10.4018/978-1-60960-491-2.ch019

Integration (ITI), was applied to several breast cancer DNA microarray datasets and allowed the extraction of a signature constituted by 119 subnetworks. All subnetworks have been stored in a relational database and linked to Gene Ontology and NCBI EntrezGene annotation databases for analysis. Exploration of annotations has shown that this set of subnetworks reflects several biological processes linked to cancer and is a good candidate for establishing a network-based signature for prediction of metastatic relapse in breast cancer.

INTRODUCTION

Since introduction of high-throughput technologies in molecular biology in the late nineties, a number of technologies for deciphering the genomic origin of several diseases has flourished. Among these, cDNA microarrays (Skena et al., 1995) have allowed measuring Gene Expression Profiles (GEP) at the genome scale and have shed light on large scale gene regulation/misregulation under varied conditions. Many diseases, including several forms of cancer [leukemia (Golub et al., 1999), colon cancer (Li et al., 2001), breast cancer (Wang et al., 2005)], diabetes (Kaestner et al., 2003), and others (Munro & Perreau. 2009) have been studied that way. Of particular interest in the context of cancer, a particularly heterogeneous disease, is the use of GEPs to either predict drug resistance (de Lavallade et al, 2010), or the metastatic recurrence, for instance in breast cancer (van de Vijver et al., 2002). Tumor microenvironment studies have allowed understanding the influence of immune system on patient outcome (Pagès et al., 2009).

There is an increasing number of controversial cases for the use of systemic adjuvant therapy due to the clinical and pathological heterogeneity of the disease to treat. In node-negative early breast cancer, most patients undergo adjuvant chemotherapy even though 70-80% of them would have survived without it (Bertucci & Birnbaum, 2008). The refinement of current prognostic histopathological methods using molecular diagnostics can also lead to increase of detection of disease subtypes that necessitate specific treatments, such as T1 breast cancer (Mook et al., 2010). In all cases,

the goal is to refine and individualize treatment and lead the way to personalized medicine for a growing number of pathologies.

In cancer, the understanding of molecular basis of metastasis is of primary importance. Several studies have attempted to obtain a molecular portrait for a large number of patients using DNA microarray analysis, performed supervised analysis and published list of genes predicting patient outcome. Two of these signatures are currently under clinical trials in breast cancer: the MIND-ACT trial, based on the 70-genes Mammaprint signature [van't Veer et al. (2002) van de Vijver et al. (2002), Bueno-de-Mesquita et al. (2007)], and the TAYLORx trial, an RT-PCR-based 21-genes OncotypeDX signature (Paik et al., 2006).

However, most prognostic signatures reported for breast cancer show very little or no overlap, and do not appear generalizable from one study to another, and this un-reproducibility was widely criticized (Chuang et al., 2007, Bertucci et al., 2008). Two studies in particular are often cited for their lack of agreement, although they addressed similar questions, which were the two breast cancer studies performed by van de Vijver et al. (2002) and Wang et al. (2005), who reported two prognosis signatures for metastatic relapse in breast cancer. Two different signatures comprising respectively 70 and 76 genes predictive of breast cancer patient outcome were reported but presented only three genes in common. Even more concerning was the study by Michiels et al. (2005) which showed that hundreds of 70-genes signatures with equal classification power can be drawn by shuffling the training and test sets in the van't Veer et al. (2002) dataset, showing the

20 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/linking-interactome-disease/52327

Related Content

Competitive Advantage of Geographical Clusters

V. Albino (2007). *Handbook of Research on Nature-Inspired Computing for Economics and Management* (pp. 317-334).

www.irma-international.org/chapter/competitive-advantage-geographical-clusters/21137

Neural Network Model for Semantic Analysis of Sanskrit Text

Smita Selot, Neeta Tripathi and A. S. Zadgaonkar (2018). *International Journal of Natural Computing Research* (pp. 1-14).

www.irma-international.org/article/neural-network-model-for-semantic-analysis-of-sanskrit-text/204880

A Comprehensive Analysis on Biodiesel Blend Model

Yarrapragada K.S.S. Rao and Bala Krishna B. (2018). *International Journal of Artificial Life Research* (pp. 25-46).

www.irma-international.org/article/a-comprehensive-analysis-on-biodiesel-blend-model/227072

Visual Tracking Using Multimodal Particle Filter

Tony Tung and Takashi Matsuyama (2014). *International Journal of Natural Computing Research* (pp. 69-84).

www.irma-international.org/article/visual-tracking-using-multimodal-particle-filter/118158

Endometrial Cancer Detection Using Pipeline Biopsies Through Machine Learning Techniques

Vemasani Varshini, Maheswari Raja and Sharath Kumar Jagannathan (2024). *Bio-Inspired Optimization Techniques in Blockchain Systems* (pp. 130-149).

www.irma-international.org/chapter/endometrial-cancer-detection-using-pipeline-biopsies-through-machine-learning-techniques/338088