

Pseudo Independent Models

Yang Xiang


University of Guelph, Canada

INTRODUCTION

Graphical models such as Bayesian networks (BNs) (Pearl, 1988) and decomposable Markov networks (DMNs) (Xiang, Wong & Cercone, 1997) have been applied widely to probabilistic reasoning in intelligent systems. Figure 1 illustrates a BN and a DMN on a trivial uncertain domain: A virus can damage computer files, and so can a power glitch. A power glitch also causes a VCR to reset. The BN in (a) has four nodes, corresponding to four binary variables taking values from {true, false}. The graph structure encodes a set of dependence and independence assumptions (e.g., that f is directly dependent on v , and p but is independent of r , once the value of p is known). Each node is associated with a conditional probability distribution conditioned on its parent nodes (e.g., $P(f|v, p)$). The joint probability distribution is the product $P(v, p, f, r) = P(f|v, p) P(r|p) P(v) P(p)$. The DMN in (b) has two groups of nodes that are maximally pair-wise connected, called *cliques*. Each clique is associated with a probability distribution (e.g., clique $\{v, p, f\}$ is assigned $P(v, p, f)$). The joint probability distribution is $P(v, p, f, r) = P(v, p, f) P(r, p) / P(p)$, where $P(p)$ can be derived from one of the clique distributions. The networks, for instance, can be used to reason about whether there are viruses in the computer system, after observations on f and r are made.

Construction of such networks by elicitation from domain experts can be very time-consuming. Automatic discovery (Neapolitan, 2004) by exhaustively testing all possible network structures is intractable. Hence, heuristic search must be used. This article examines a class of graphical models that cannot be discovered using the common heuristics.

Figure 1. (a) a trivial example BN; (b) a corresponding DMN



BACKGROUND

Let V be a set of n discrete variables x_1, \dots, x_n (in what follows, we will focus on finite, discrete variables). Each variable x_i has a finite space $S_i = \{x_{i,1}, x_{i,2}, \dots, x_{i,D}\}$ of cardinality D_i . When there is no confusion, we write $x_{i,j}$ as x_{ij} for simplicity. The space of a set V of variables is defined by the Cartesian product of the spaces of all variables in V , that is, $S_V = S_1 \times \dots \times S_n$ (or Π^S). Thus, S_V contains the tuples made of all possible combinations of values of the variables in V . Each tuple is called a configuration of V , denoted by $\mathbf{v} = (x_1, \dots, x_n)$.

Let $P(x_i)$ denote the probability function over x_i and $P(x_{ij})$ denote the probability value $P(x_i = x_{ij})$. A probabilistic domain model (PDM) M over V defines the probability values of every configuration for every subset $A \subseteq V$. Let $P(V)$ or $P(x_1, \dots, x_n)$ denote the joint probability distribution (JPD) function over x_1, \dots, x_n and $P(x_{1j_1}, \dots, x_{nj_n})$ denote the probability value of a configuration $(x_{1j_1}, \dots, x_{nj_n})$. We refer to the function $P(A)$ over $A \subseteq V$ as the *marginal distribution* over A and $P(x_i)$ as the *marginal distribution* of x_i . We refer to $P(x_{1j_1}, \dots, x_{nj_n})$ as a *joint parameter* and $P(x_{ij})$ as a *marginal parameter* of the corresponding PDM over V .

For any three disjoint subsets of variables W, U and Z in V , subsets W and U are called *conditionally independent* given Z , if

$$P(W | U, Z) = P(W | Z)$$

for all possible values in W, U and Z such that $P(U, Z) > 0$. Conditional independence signifies the dependence mediated by Z . This allows the dependence among $W \cup U \cup Z$ to be modeled over subsets $W \cup Z$ and $U \cup Z$ separately. Conditional independence is the key property explored through graphical models.

Subsets W and U are said to be *marginally independent* (sometimes referred to as *unconditionally independent*) if

$$P(W | U) = P(W)$$

for all possible values W and U such that $P(U) > 0$. When two subsets of variables are marginally independent, there is no dependence between them. Hence, each subset

can be modeled independently without losing information.

If each variable x_i in a subset A is marginally independent of $A \setminus \{x_i\}$, the variables in A are said to be *marginally independent*. The following proposition reveals a useful property called *factorization* when this is the case.

- **Proposition 1:** If each variable x_i in a subset A is marginally independent of $A \setminus \{x_i\}$ then

$$P(A) = \prod_{x_i \in A} P(x_i)$$

Variables in a subset A are called *generally dependent*, if $P(B | A \setminus B) \neq P(B)$ for every proper subset $B \subset A$. If a subset of variables is generally dependent, its proper subsets cannot be modeled independently without losing information. A generally dependent subset of variables, however, may display conditional independence within the subset. For example, consider $A = \{x_1, x_2, x_3\}$. If $P(x_1, x_2 | x_3) = P(x_1, x_2)$, i.e., $\{x_1, x_2\}$ and x_3 are marginally independent, then A is *not* generally dependent. On the other hand, if

$$P(x_1, x_2 | x_3) \neq P(x_1, x_2), P(x_2, x_3 | x_1) \neq P(x_2, x_3), P(x_3, x_1 | x_2) \neq P(x_3, x_1),$$

then A is generally dependent.

Variables in A are *collectively dependent* if, for each proper subset $B \subset A$, there exists no proper subset $C \subset A \setminus B$ that satisfies $P(B | A \setminus B) = P(B | C)$. Collective dependence prevents conditional independence and modeling through proper subsets of variables. Table 1 shows the JPD over a set of variables $V = \{x_1, x_2, x_3, x_4\}$. The four variables are collectively dependent; for example,

$$P(x_{1,1}, x_{2,0}, x_{3,1}, x_{4,0}) = 0.257$$

and

$$P(x_{1,1}, x_{2,0}, x_{3,1}) = P(x_{1,1}, x_{2,0}, x_{4,0}) = P(x_{1,1}, x_{3,0}, x_{4,0}) = 0.3.$$

MAIN THRUST

Pseudo-Independent (PI) Models

A *pseudo-independent* (PI) model is a PDM where proper subsets of a set of collectively dependent variables display marginal independence (Xiang, Wong & Cercone, 1997). The basic PI model is a full PI model:

- **Definition 2 (Full PI model):** A PDM over a set $V (|V| \geq 3)$ of variables is a *full PI model*, if the following properties (called axioms of full PI models) hold:

(S_I) Variables in each proper subset of V are marginally independent.

(S_{II}) Variables in V are collectively dependent.

Table 1 shows the JPD of a binary full PI model, where $V = \{x_1, x_2, x_3, x_4\}$. Its marginal parameters are

$$P(x_{1,0}) = 0.7, P(x_{2,0}) = 0.6, P(x_{3,0}) = 0.35, P(x_{4,0}) = 0.45.$$

Any subset of three variables are marginally independent; for example,

$$P(x_{1,1}, x_{2,0}, x_{3,1}) = P(x_{1,1}) P(x_{2,0}) P(x_{3,1}) = 0.117.$$

The four variables are collectively dependent as explained previously.

Table 2 is the JPD of the color model given earlier, where $V = \{x_1, x_2, x_3\}$. The marginal independence can be verified by

$$P(x_1 = \text{red}) = P(x_2 = \text{red}) = P(x_3 = \text{red}) = 0.5, \\ P(x_1 = \text{red} \setminus x_2) = P(x_1 = \text{red} \setminus x_3) = P(x_2 = \text{red} \setminus x_3) = 0.5,$$

and the collective dependence can be seen from $P(x_1 = \text{red} \setminus x_2 = \text{red}, x_3 = \text{red}) = 1$.

By relaxing condition (S_I) on marginal independence, full PI models are generalized into partial PI models, which are defined through marginally independent partition (Xiang, Hu, Cercone & Hamilton, 2000) introduced in the following:

- **Definition 3 (Marginally Independent Partition):** Let $V (|V| \geq 3)$ be a set of variables, and $B = \{B^1, \dots, B^m\} (m \geq 2)$ be a partition of V . B is a marginally independent partition if, for every subset $A = \{x_i^k | x_i^k \in B^k, k = 1, \dots, m\}$, variables in A are marginally independent. Each block B^k is called a *marginally independent block*.

Intuitively, a marginally independent partition groups variables in V into m blocks. If one forms a subset A by taking one element from each block, then variables in A are marginally independent. Unlike full PI models, in a partial PI model, it is not necessary that every proper subset is marginally independent. Instead, that requirement is replaced with the marginally independent partition.

- **Definition 4 (Partial PI Model):** A PDM over a set $V (|V| \geq 3)$ of variables is a partial PI model, if the following properties (called axioms of partial PI models) hold:

(S_I') V can be partitioned into two or more marginally independent blocks.

4 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/chapter/pseudo-independent-models/10730

Related Content

Retrieving Medical Records Using Bayesian Networks

Luis M. de Campos, Juan M. Fernandez-Luna and Juan F. Huete (2005). *Encyclopedia of Data Warehousing and Mining* (pp. 960-964).

www.irma-international.org/chapter/retrieving-medical-records-using-bayesian/10735

Enhancing Web Search through Web Structure Mining

Ji-Rong Wen (2005). *Encyclopedia of Data Warehousing and Mining* (pp. 443-447).

www.irma-international.org/chapter/enhancing-web-search-through-web/10638

Database Sampling for Data Mining

Patricia E.N. Lutu (2005). *Encyclopedia of Data Warehousing and Mining* (pp. 344-348).

www.irma-international.org/chapter/database-sampling-data-mining/10620

Managing Late Measurements in Data Warehouses

Matteo Golfarelli and Stefano Rizzi (2008). *Data Warehousing and Mining: Concepts, Methodologies, Tools, and Applications* (pp. 738-754).

www.irma-international.org/chapter/managing-late-measurements-data-warehouses/7673

Data Warehousing and Data Mining Lessons for EC Companies

Neerja Sethi and Vijay Sethi (2008). *Data Warehousing and Mining: Concepts, Methodologies, Tools, and Applications* (pp. 381-388).

www.irma-international.org/chapter/data-warehousing-data-mining-lessons/7653