

Chapter 8

Cluster Origin of Solvation Features of C–Nanostructures in Organic Solvents

Francisco Torrens

Universitat de València, Spain

Gloria Castellano

Universidad Católica de Valencia, Spain

ABSTRACT

The existence of fullerenes, Single-Wall Carbon Nanocones (SWNCs), especially Nanohorns (SWNHs), Single-Wall Carbon Nanotube (SWNT) (CNT) (NT), NT-Fullerene Bud (NT-BUD), Nanographene (GR) and GR-Fullerene Bud (GR-BUD) in cluster form is discussed in organic solvents. Theories are developed based on columnlet, bundlet and droplet models describing size-distribution functions. The phenomena present a unified explanation in the columnlet model in which free energy of cluster-involved GR comes from its volume, proportional to number of molecules n in cluster. Columnlet model enables describing distribution function of GR stacks by size. From geometrical considerations, columnlet (GR/GR-BUD), bundlet (SWNT/NT-BUD) and droplet (fullerene) models predict dissimilar behaviours. Interaction-energy parameters are derived from C_{60} . An NT-BUD behaviour or further is expected. Solubility decays with temperature result smaller for GR/GR-BUD than SWNT/NT-BUD than C_{60} in agreement with lesser numbers of units in clusters. Discrepancy between experimental data of the heat of solution of fullerenes, CNT/NT-BUDs and GR/GR-BUDs is ascribed to the sharp concentration dependence of the heat of solution. Diffusion coefficient drops with temperature result greater for GR/GR-BUD than SWNT/NT-BUD than C_{60} corresponding to lesser number of units in clusters. Aggregates $(C_{60})_{13}$, SWNT/NT-BUD₇ and GR/GR-BUD₃ are representative of droplet, bundlet and columnlet models.


DOI: 10.4018/978-1-5225-0248-7.ch008

INTRODUCTION

Interest in nanoparticles (NPs) arises from the shape-dependent physical properties of materials at the nanoscale (Faraday, 1857; Murphy *et al.*, 2010). Occurrence of single-wall carbon nanocones (SWNCs) was used to investigate nucleation and growth of curved C-nanostructures (NSs) suggesting pentagon role. When a pentagon is introduced into a graphitic sheet nanographene (GR) (Figure 1d) *via* extraction of a 60° sector from the sheet one forms a cone leaf. Pentagons presence in an SWNC apex is analogue of their occurrence in single-wall C-nanotube (NT) (CNT) (SWNT) tip topology (*cf.* Figure 1b). Terminations of SWNTs attracted interest once Tamura & Tsukada (1995) theoretically predicted peculiar electronic states related to GR topological defects. Kim *et al.* (1999) observed resonant peaks in density of states (DOS) in SWNTs and Carroll *et al.* (1997), in multiple-wall (MNTs) C-nanotubes (MWNTs).

The SWNCs with discrete opening angles (apices, θ) of 19°, 39°, 60°, 85° and 113° of cone (*cf.* Figure 2) were observed in a C-sample generated by hydrocarbon (HC) pyrolysis (Krishnan *et al.*, 1997), which was explained by a cone-wall model composed of wrapped GR sheets where geometrical requirement for seamless connection accounted for semidiscrete character and absolute values of cone angle. Total

Figure 1. Arrangement of C-nanostructures: (a) C_{60} ; (b) SWNT; (c) NT-BUD; (d) GR; (e) GR-BUD



103 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/cluster-origin-of-solvation-features-of-c-nanostructures-in-organic-solvents/150020

Related Content

Adapting Education to the Needs of Pharmacists and Pharmaceutical Scientists

Bruno Sepodes, João Pedro Rocha and Maria-Eduardo Figueira (2021). *Pedagogies for Pharmacy Curricula* (pp. 20-42).

www.irma-international.org/chapter/adapting-education-to-the-needs-of-pharmacists-and-pharmaceutical-scientists/269627

Protein Ligand Interaction Fingerprints

Ali HajiEbrahimi, Hamidreza Ghafouri, Mohsen Ranjbar and Amirhossein Sakhteman (2016). *Methods and Algorithms for Molecular Docking-Based Drug Design and Discovery* (pp. 128-147).

www.irma-international.org/chapter/protein-ligand-interaction-fingerprints/151885

Computational Approaches for the Discovery of Novel Hepatitis C Virus NS3/4A and NS5B Inhibitors

Khac-Minh Thai, Quoc-Hiep Dong, Thi-Thanh-Lan Nguyen, Duy-Phong Le, Minh-Tri Le and Thanh-Dao Tran (2015). *Quantitative Structure-Activity Relationships in Drug Design, Predictive Toxicology, and Risk Assessment* (pp. 318-353).

www.irma-international.org/chapter/computational-approaches-for-the-discovery-of-novel-hepatitis-c-virus-ns34a-and-ns5b-inhibitors/124474

Phytochemistry, Ethnobotany, Biogenesis, and Pharmacological Wonders of Cumin Seeds

Ammara Chand, Saima Ali and Saikh Mohammad Wabaidur (2024). *Therapeutic and Pharmacological Applications of Ethnobotany* (pp. 128-154).

www.irma-international.org/chapter/phytochemistry-ethnobotany-biogenesis-and-pharmacological-wonders-of-cumin-seeds/344959

Therapeutic Properties of Syzygium cumini (Jamun) and Tinospora cordifolia (Giloy) Against Various Lethal Diseases

Ankita Gautam (2020). *Advanced Pharmacological Uses of Medicinal Plants and Natural Products* (pp. 316-337).

www.irma-international.org/chapter/therapeutic-properties-of-syzygium-cumini-jamun-and-tinospora-cordifolia-giloy-against-various-lethal-diseases/252952