

# Chapter 15

## Prediction Changes for Nonstationary Multi–Temporal Satellite Images using HMM

**Ali Ben Abbes**  
*ENSI, Tunisia*

**Imed Riadh Farah**  
*ISAMM, Tunisia*

### ABSTRACT

*Due to the growing advances in their temporal, spatial, and spectral resolutions, remotely sensed data continues to provide tools for a wide variety of environmental applications. This chapter presents the benefits and difficulties of Multi-Temporal Satellite Image (MTSI) for land use. Predicting land use changes using remote sensing is an area of interest that has been attracting increasing attention. Land use analysis from high temporal resolution remotely sensed images is important to promote better decisions for sustainable management land cover. The purpose of this book chapter is to review the background of using Hidden Markov Model (HMM) in land use change prediction, to discuss the difference on modeling using stationary as well as non-stationary data and to provide examples of both case studies (e.g. vegetation monitoring, urban growth).*

### INTRODUCTION

Recent years have seen growing interest in technological evolution of remote sensing sensors and satellites which provides a huge amount of free valuable earth observation data (Lillesand et al., 2014; Reiche et al., 2015). Therefore, a crucial importance is given to develop and to improve innovative tools in order to make the most of its spatio-temporal informational content (Senthilnath et al., 2014; Wondrade, et al., 2014; Badreldin & Goossens, 2014). The accessibility of remotely sensed images and growing advances in their temporal, spatial, and spectral resolutions continues to provide tools for detecting land use changes at different scales. Multi-Temporal Satellite Images (MTSI) analysis from high temporal resolution remotely sensed images is important to promote better decisions for sustainable management

DOI: 10.4018/978-1-5225-0937-0.ch015

of land use. The free of cost of long MTSI of, for example, Landsat and MODIS makes the development and analysis of land use techniques a hot topic in the field of remote sensing. Currently, much attention has been given to land use in order to improve our understanding of the causes and consequences of these changes. Notably, to analyze the possible scenarios of land use change in the future.


This chapter answers this issue to benefit from these informations extracted from MTSI to detect land use changes. Land use is a major issue due to its practical uses in various applications, such as: deforestation, damage assessment, disasters monitoring, urban expansion, planning, and land management. Accurate and up-to-date land use data is necessary to understand and assess the change. This issue was addressed in several research papers and tackled from different points of view (Coppin et Bauer, 1996; Veldkam & Lambin, 2001Coppin, et al., 2004Badreldin & Goossens, 2014; Halmy et al., 2015). Consequently, two significant challenges stand out. First, methods must allow change detection for nonstationarity in MTSI. Secondly, methods must build a prediction system to predict future state of land use. The aim of land use changes prediction is to predict and analyze the future scenario for land use. Predictive models comprise a variety of methods based on statistics and data mining that analyze current and historical data to make a prediction about future events. Generally, predictive analytics can help experts to understand land use change, to identify unpredicted opportunities and to anticipate changes before they happen. Over the last decades, a variety of methods have been developed for change prediction using remotely sensed images such as neural networks (Li & Yeh, 2002; Basse et al., 2014), decision trees (Bui et al., 2015; Bouatay et al., 2014), Markov models (Viovy & Saint, 1994; Salberg, et al., 2011; Shen et al., 2013; Leite et al., 2014), multi-agent systems (Parker et al., 2003; Hosseinali & Alesheikh, 2014), statistical models (Faghih-Imani et al., 2014; Rosà, et al., 2014) and evolution model (Boulangeat et al., 2014 ; Riordan and Rundel, 2014).

The most generic schema of land use prediction (as shown in Figure 1) (Halmy et al., 2015), considers as input a sequence of satellite images, which describe a geographical area (e.g. vegetation, urban) at different times. Comprises broadly speaking:

1. Pre-process images,
2. Extract interest area,
3. Calculate and normalize attributes and
4. Build prediction model.

This chapter addresses the use of Hidden Markov Model (HMM) in prediction of land use change. In fact, HMMs have been widely used in change prediction in MTSI under stationarity hypothesis. Two problems in implementation of predictive analytics give off the following items:

*Figure 1. Generic system for land use prediction changes using MTSI*



18 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

[www.igi-global.com/chapter/prediction-changes-for-nonstationary-multi-temporal-satellite-images-using-hmm/169997](http://www.igi-global.com/chapter/prediction-changes-for-nonstationary-multi-temporal-satellite-images-using-hmm/169997)

## Related Content

---

### Geographical Information Systems in Modern Citizen Science

Laia Subirats, Joana Simoes and Alexander Steblin (2019). *Geospatial Intelligence: Concepts, Methodologies, Tools, and Applications* (pp. 172-201).

[www.irma-international.org/chapter/geographical-information-systems-in-modern-citizen-science/222898](http://www.irma-international.org/chapter/geographical-information-systems-in-modern-citizen-science/222898)

### Bacterial Source Tracking of Nonpoint Source Pollution Using GIS and DNA Fingerprinting Technologies

M. E. Folkoff, E. A. Venso, D. W. Harris, M. F. Frana and M. S. Scott (2003). *Geographic Information Systems and Health Applications* (pp. 208-223).

[www.irma-international.org/chapter/bacterial-source-tracking-nonpoint-source/18843](http://www.irma-international.org/chapter/bacterial-source-tracking-nonpoint-source/18843)

### Application of Data Fusion for Uncertainty and Sensitivity Analysis of Water Quality in the Shenandoah River

Mbongowo Joseph Mbuh (2019). *Geospatial Intelligence: Concepts, Methodologies, Tools, and Applications* (pp. 1383-1410).

[www.irma-international.org/chapter/application-of-data-fusion-for-uncertainty-and-sensitivity-analysis-of-water-quality-in-the-shenandoah-river/222953](http://www.irma-international.org/chapter/application-of-data-fusion-for-uncertainty-and-sensitivity-analysis-of-water-quality-in-the-shenandoah-river/222953)

### Towards the Integration of Trajectory Information Sources for Semantic Conflicts Detection Purpose: A Trajectory Ontology Based Approach

Marwa Manaa and Akaichi Jalel (2017). *Handbook of Research on Geographic Information Systems Applications and Advancements* (pp. 488-519).

[www.irma-international.org/chapter/towards-the-integration-of-trajectory-information-sources-for-semantic-conflicts-detection-purpose/170002](http://www.irma-international.org/chapter/towards-the-integration-of-trajectory-information-sources-for-semantic-conflicts-detection-purpose/170002)

### Spatiotemporal Network Analysis and Visualization

Judith Gelernter and Kathleen M. Carley (2015). *International Journal of Applied Geospatial Research* (pp. 77-97).

[www.irma-international.org/article/spatiotemporal-network-analysis-and-visualization/122363](http://www.irma-international.org/article/spatiotemporal-network-analysis-and-visualization/122363)