

# Chapter 8

## Land Use – Terrain Correlations in the Piedmont Tract of Eastern India: A Case Study of the Dulung River Basin

**Ashis Sarkar**

*Chandernagore Government College, India*

**Priyank Pravin Patel**

*Presidency University, India*

### **ABSTRACT**

*The Dulung River flows across West Bengal and Jharkhand in India. The geographical variables present within the basin area are categorised into groups like Physiographic, Morphometric and Land Use-Land Cover (LULC) attributes. These facets are mapped and overlain in a GIS environment and correlations drawn between them. Factor Scores obtained through Principal Component Analysis are further compared and correlated. The different variables are fused to obtain a comprehensive grouping of the above three facets that is reflective of the overall terrain attributes and its overlying LULC classes. Through this, within the Dulung River Basin, three broad Physiographic-Soil-Land Use Units (PSLUs) are identified, which comprise of the structural ridges and residual hills, piedmont plains and floodplains. For further insight into existing LULC-landform relations, select villages across the basin landscape are examined in detail. The relations derived help in suggesting possible land management practices in this region.*

DOI: 10.4018/978-1-5225-1814-3.ch008

## **INTRODUCTION**

Land use is a dynamic phenomenon and needs continuous monitoring (Bhattacharya, 1983). Though it is mainly controlled by climatic, pedological, socioeconomic and political conditions yet geology and geomorphology too play important roles since a physical landscape is inherently constituted by its underlying lithology and structure. Depending on their occurrence in a particular climatic region, individual landforms give rise to particular soil types and certain landform-soil associations are often more suitable for specific land uses. The terrain configuration, affecting the soil and surface drainage, is borne of the sub-surface geology, surficial landforms and their slopes. Morphometric parameters, after enumeration, could be thus used in many watershed and LULC planning applications. As such, studies which integrate landform analysis with soil, hydrology and land use and land cover (LULC) facets, should ideally form the basis of the land resource evaluation of a region (Subramanian, 1978, 1981).

LULC study of an area provides information about its present land use status and helps form a baseline study for sustainable development of that locale (Krishna, Westinga, & Huizing, 1999). This requires current and archival LULC data of the area to track temporal change patterns (Chaurasia et. al., 1996), specifically information on existing land use and the distribution of settlements, forests, agricultural and barren lands (Christian & Stewart, 1953; Natarajan, Gajbe, & Manchanda, 1986; Shankarnarayan & Sen, 1977). LULC changes are linked to the intersection of natural and human influences on environmental regimes. Biosphere changes and biogeochemical cycle alterations are triggered by heterogeneous changes in land use and manifested further through their perpetuation (Turner, 1995).

Drainage basins have always had a special relevance in geomorphology (Doornkamp & King, 1971; Strahler, 1957) and watersheds are widely adopted in most countries as natural units of ecosystem planning and development (FAO, 1977, 1987; Honore, 1999; Moore, Grayson, & Ladson, 1977). The watershed is a geo-hydrological unit of area drained to a common point and considered as an ideal unit for analysis and management of natural resources and environmental planning in any ridge-to-valley treatment, with its appropriateness in regional planning quite evident (Easter, Stow, & Jenson, 1986; Gregory & Walling, 1978; Saha & Barrow, 1981). In India too, many developmental programmes are based on watershed management (Dhruvanarayana, Sastry, & Patnaik, 1990). Furthermore, India's agrarian-cum-rural economy (on which depend over two hundred million farmers) is extremely sensitive to land and water management and even if the country's irrigation potential is maximised, about 50% of the net cultivated area will continue to depend on rainfall (ISRO/RRSSC, 2009). Much focus thus needs to be given to

44 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: [www.igi-global.com/chapter/land-use---terrain-correlations-in-the-piedmont-tract-of-eastern-india/172711](http://www.igi-global.com/chapter/land-use---terrain-correlations-in-the-piedmont-tract-of-eastern-india/172711)

## Related Content

---

### The Circular Economy, Big Data Analytics, and the Transformation of Urban Slums in Sub-Saharan Africa

Darrold Laurence Cordes and Gregory Morrison (2023). *International Journal of Smart Sensor Technologies and Applications* (pp. 1-27).

[www.irma-international.org/article/the-circular-economy-big-data-analytics-and-the-transformation-of-urban-slums-in-sub-saharan-africa/319720](http://www.irma-international.org/article/the-circular-economy-big-data-analytics-and-the-transformation-of-urban-slums-in-sub-saharan-africa/319720)

### Blockchain Hyperledger Sawtooth Enabled Digital Forensics Chain of Custody (CoC) A Short Report

(2022). *The International Journal of Imaging and Sensing Technologies and Applications* (pp. 0-0).

[www.irma-international.org/article//306655](http://www.irma-international.org/article//306655)

### Diameter-Aggregation Delay Tradeoff for Data Gathering Trees in Wireless Sensor Networks

Natarajan Meghanathan (2020). *Sensor Technology: Concepts, Methodologies, Tools, and Applications* (pp. 335-350).

[www.irma-international.org/chapter/diameter-aggregation-delay-tradeoff-for-data-gathering-trees-in-wireless-sensor-networks/249570](http://www.irma-international.org/chapter/diameter-aggregation-delay-tradeoff-for-data-gathering-trees-in-wireless-sensor-networks/249570)

### Visual Sensor Network Technology and Its Applications

Li-minn Ang and Kah Phooi Seng (2012). *Visual Information Processing in Wireless Sensor Networks: Technology, Trends and Applications* (pp. 1-19).

[www.irma-international.org/chapter/visual-sensor-network-technology-its/59750](http://www.irma-international.org/chapter/visual-sensor-network-technology-its/59750)

### Designing Mobile Learning Smart Education System Architecture for Big Data Management Using Fog Computing Technology

Muhammad Adnan Kaim Khani, Abdullah Ayub Khan, Allah Bachayo Brohi and Zaffar Ahmed Shaikh (2022). *The International Journal of Imaging and Sensing Technologies and Applications* (pp. 1-23).

[www.irma-international.org/article/designing-mobile-learning-smart-education-system-architecture-for-big-data-management-using-fog-computing-technology/306653](http://www.irma-international.org/article/designing-mobile-learning-smart-education-system-architecture-for-big-data-management-using-fog-computing-technology/306653)