

Chapter 7

Non-Ionic Surfactant Vesicles (Niosomes) as New Drug Delivery Systems

Abbas Pardakhty
Kerman University of Medical Sciences, Iran

ABSTRACT

Lipid vesicular systems composed of hydrated amphiphiles with or without bilayer inducing agents such as cholesterol. On the basis of used amphiphilic molecule different nomenclature are used as liposomes, ufasomes and niosomes. Nonionic surfactants with mono-, di- or trialkyl chains form niosomes which are lipid vesicles with more chemical stability in comparison with phospholipids of liposomes. Both hydrophobic and hydrophilic chemicals can be encapsulated in niosomes as a new drug delivery system. This drug carrier system could have administered via injection, oral, pulmonary, vaginal, rectal, ophthalmic, nasal or transdermal routes with penetration enhancing potential. This chapter presents a detailed explain about niosome forming components, methods of preparation and routes of administration. Many examples for drug delivery potential of niosomes are also available in this review. Vaccine adjuvant and genetic substances vector capabilities are not given here.

INTRODUCTION

There are versatile lists of amphiphilic compounds for production of lipid vesicles which on the basis of used amphiphiles or structural compounds, different nomenclature has been used such as liposomes (Kaminskas et al. 2012), niosomes (Moazeni et al. 2010; Akbari et al. 2013), ufasomes (Patel et al. 2011), ethosomes (Liu et al. 2012), eliposomes (Lattin et al. 2012) and so on. In lipid vesicular systems, polar headgroups of amphiphilic molecules are oriented into aqueous compartments and their lipophilic parts form bilayer membranes. Niosomes are bilayer vesicles composed mainly of hydrated non-ionic surfactants in addition to, in many cases, cholesterol (CHOL) or its derivatives (Pardakhty and Moazeni 2013). Cosmetic industry was the place for the first account of niosome production (Handjani-Vila et Al. 1979) after which a large number of niosome applications in drug, vaccine and gene delivery have been

DOI: 10.4018/978-1-5225-1762-7.ch007


explored. While most niosomes are in the nano or sub-micron (colloidal) size range, not many authors used the “nanoniosome” or “nanovesicle” titles in their published articles due to introduction the new nanotechnology related phrases during the few recent years. Niosomes are capable to encapsulate both hydrophilic and lipophilic substances which the first ones usually are entrapped in vesicular aqueous core or adsorbed on the bilayer surfaces, and lyophobic molecules are encapsulated by their partitioning into the lipophilic domain of the bilayers. Incorporation of charged molecules into non-ionic surfactant vesicles forms charged or hybrid niosomes (Sennato et al. 2008). Vesicles can be also prepared from the mixed cationic and anionic surfactant systems such as cetyltrimethylammonium bromide and sodium lauryl sulfate (CTAB-SLS) (Aiello et al. 2010). These types of vesicles are presented as “cat-anionic” or “catanionic” vesicles. Catanionic vesicles can also be formed from electrostatic interaction between an ionized drug, such as diclofenac (Zhao et al. 2013) or alprenolol (Dew et al. 2012) and a positively/negatively charged surfactant.

Non-ionic surfactants have more chemical stability against oxidation and are less thermo-labile in comparison to phospholipids, the main constituent of liposomes (Fang et al. 2001; Tarekegn et al. 2010) thus less significant care in handling and storage are needed. Furthermore, greater versatility and lower cost make this type of vesicles more attractive in drug, gene and vaccine delivery (Florence 1993). From the pharmaceutical manufacturing stand of view the superiority of niosomes is simple methods required for industrialized and large-scale production of them without the use of pharmaceutically unacceptable solvents (Sahin 2007). In this chapter the most important aspects of niosomes and drug delivery potentials will be reviewed. Application of niosomes as gene or vaccine delivery tools will not discussed in this chapter.

Amphiphiles (Surfactants)

Following the application of some forms of energy such as mechanical or heating, the formation of liposomes is a self-assembly process due to high interfacial tension between aqueous medium and the lipophilic alkyl chain(s) resulted in the association of non-ionic surfactant monomers into vesicles (Uchegbu and Vyas 1998). Concurrently, the hydrophilic headgroups of amphiphilic molecules make water mediated interactions counter the previous formed force, finally result in bilayer formation. The

Figure 1. α,ω -trioxyethylene-bis(sodium 2-dodecyloxy-propylenesulfonate), an anionic gemini surfactant, used for preparation of niosomes
Muzzalupo et al. 2008.



29 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/non-ionic-surfactant-vesicles-niosomes-as-new-drug-delivery-systems/174125

Related Content

Quantitative Structure-Activity/Property/Toxicity Relationships through Conceptual Density Functional Theory-Based Reactivity Descriptors

Sudip Pan, Ashutosh Kumar Gupta, Venkatesan Subramanian and Pratim K. Chattaraj (2017).

Pharmaceutical Sciences: Breakthroughs in Research and Practice (pp. 1517-1572).

www.irma-international.org/chapter/quantitative-structure-activitypropertytoxicity-relationships-through-conceptual-density-functional-theory-based-reactivity-descriptors/174180

Ethnobotanical, Therapeutic, and Pharmacological Applications of Bakuchi (*Psoralea corylifolia* L.) Plant

Sandipan Babasaheb Jige (2024). *Therapeutic and Pharmacological Applications of Ethnobotany* (pp. 23-37).

www.irma-international.org/chapter/ethnobotanical-therapeutic-and-pharmacological-applications-of-bakuchi-psoralea-corylifolia-l-plant/344955

Enzymatic Research Having Pharmaceutical Significance

Ishan H. Ravaland Arvind Kumar Singh Chandel (2018). *Research Advancements in Pharmaceutical, Nutritional, and Industrial Enzymology* (pp. 141-158).

www.irma-international.org/chapter/enzymatic-research-having-pharmaceutical-significance/203814

Cyanobacterial Toxins in Water Sources and Their Impacts on Human Health

Zakaria Mohamed (2017). *Pharmaceutical Sciences: Breakthroughs in Research and Practice* (pp. 1428-1456).

www.irma-international.org/chapter/cyanobacterial-toxins-in-water-sources-and-their-impacts-on-human-health/174176

Implementation of Nanoparticles in Cancer Therapy

Ece Bayir, Eyup Bilgi and Aylin Sendemir Urkmez (2017). *Pharmaceutical Sciences: Breakthroughs in Research and Practice* (pp. 1212-1257).

www.irma-international.org/chapter/implementation-of-nanoparticles-in-cancer-therapy/174168