

Chapter 9

An Artificial Intelligence Approach to Thrombophilia Risk

João Vilhena

Universidade de Évora, Portugal

Henrique Vicente

Universidade de Évora, Portugal

M. Rosário Martins

Universidade de Évora, Portugal

José Grañeda

Hospital do Espírito Santo de Évora, Portugal

Filomena Caldeira

Hospital do Espírito Santo de Évora, Portugal

Rodrigo Gusmão

Hospital do Espírito Santo de Évora, Portugal

João Neves

Drs. Nicolas & Asp, UAE

José Neves

Universidade do Minho, Portugal

ABSTRACT

Thrombophilia stands for a genetic or an acquired tendency to hypercoagulable states, frequently as venous thrombosis. Venous thromboembolism, represented mainly by deep venous thrombosis and pulmonary embolism, is often a chronic illness, associated with high morbidity and mortality. Therefore, it is crucial to identify the cause of the disease, the most appropriate treatment, the length of treatment or prevent a thrombotic recurrence. This work will focus on the development of a diagnosis decision support system in terms of a formal agenda built on a Logic Programming approach to knowledge representation and reasoning, complemented with a computational framework based on Artificial Neural Networks. The proposed model has been quite accurate in the assessment of thrombophilia predisposition (accuracy close to 95%). Furthermore, the model classified properly the patients that really presented the pathology, as well as classifying the disease absence (sensitivity and specificity higher than 95%).

DOI: 10.4018/978-1-5225-7122-3.ch009

INTRODUCTION

Thrombophilia is a genetic or acquired tendency to hypercoagulable states that increase the risk of arterial and often, venous thrombosis. Venous ThromboEmbolism (VTE) is represented by two main clinical events: Deep Venous Thrombosis (DVT) and Pulmonary Embolism (PE) which often constitute a unique clinical picture in which PE follows DVT (Previtaly, Bucciarelli, Passamonti, & Martinelli, 2011). VTE is a multifactorial disease, often asymptomatic and associated with high morbidity and mortality (East & Wakefield, 2010).

The two most common factors for inherited (genetic) thrombophilias are the V Leiden mutation and the prothrombin G20210A gene mutation. Other mutations or polymorphisms associated with increased risk of thrombosis are methylenetetra-hydrofolate reductase 677C (Lim & Moll, 2015). People can have one abnormal gene (heterozygous state) or two abnormal genes (homozygous state). Less common genetic thrombophilia include deficiencies of the blood clotting proteins called protein C, protein S, and antithrombin (Lim & Moll, 2015; Previtaly et al., 2011).

Secondary hypercoagulable states are a frequently acquired VTE in patients with underlying systemic diseases or clinical conditions, known to be associated with an increased risk of thrombosis (e.g., malignancy, pregnancy, and use of oral contraceptives, myeloproliferative disorders, hyperlipidemia, diabetes mellitus and abnormalities of blood vessels). The most common acquired thrombophilia is antiphospholipid antibody syndrome (Previtaly et al., 2011).

The incidence of VTE is estimated at 56-160/100.000 people/year (East & Wakefield, 2010) and is strongly age-dependent, rising nearly 1% per year in old age (Rosendaal, 2005). Studies show that about 70% of patients presenting the first episode of VTE are over 60 years old, and the rate of recurrence is higher when the first episode of VTE occurs before 60 years old (Pernod et al., 2009). Moreover, a non-O blood group is associated with an increased risk of proximal deep vein thrombosis of the lower limbs with or without pulmonary embolism. The addition of inherited thrombophilia increases the thrombotic risk conferred by non-O group alone by almost 3-fold (Spiezia et al., 2013). Environmental or acquired VTE risk factors also include previous immobility, surgery, obesity, smoking habits and pregnancy or postpartum status (Anderson & Spencer, 2003; Enga et al., 2012; Goldhaber, 2010). Furthermore, the long distance travel is often associated with an up to 4-fold increased risk of VTE compared to non-travelers (Bartholomew, Schaffer, & McCormick, 2011).

Follow-up of patients for prolonged periods after an initial DVT or PE has revealed that VTE often is a chronic illness requiring lifelong prevention strategies. Thus, it is crucial to identify the cause, the most appropriate treatment, how long the treatment should be or how to prevent a thrombotic recurrence (Sinescu, Hostiuc, & Bartos, 2011). In the case of patients with a high risk of clinical complications preventive actions should be taken. The screening for VTE is mandatory in three high-risk groups, namely women who are prescribed oral estrogen preparations, pregnancy morbidity and patients undergoing major orthopedic surgery (Wu et al., 2006). Thus, more studies are necessary to reach a correct identification of factors associated with these diseases in order to assess the individual risk of thrombosis, and promote more targeted prophylactic and therapeutic alternatives.

On the one hand, in this study the complex pathophysiologic features of these hypercoagulable states (i.e., genetic, environmental and acquired risk factors) are discussed, being the problem tackled with Artificial Intelligence (AI) based methodologies and techniques for problem-solving (Chandrasekaran, 1983). Indeed, this work will focus on the development of an AI grounded Decision Support System aiming at the early diagnosis of thrombophilia, and also to signaling of patients with hypercoagulable

20 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/an-artificial-intelligence-approach-to-thrombophilia-risk/213344

Related Content

Prevalence of Cardiovascular Risk Factors in the Elderly Population of Alentejo: Integrative Review

Maria do Céu Mendes Pinto Marques, Ana Filipa Pereira Vaz, Ana Sofia Emídio Cardoso Leite, Cláudia Sofia Araújo, Cláudia Roque Condeçoand José Ribeiro de Deus (2020). *Handbook of Research on Health Systems and Organizations for an Aging Society* (pp. 185-195).

www.irma-international.org/chapter/prevalence-of-cardiovascular-risk-factors-in-the-elderly-population-of-alentejo/238280

Medicinal Plants for the Treatment of Liver Cancer

Bui Thanh Tung, Dang Kim Nganand Do Thi Hong Khanh (2022). *Handbook of Research on Natural Products and Their Bioactive Compounds as Cancer Therapeutics* (pp. 232-248).

www.irma-international.org/chapter/medicinal-plants-for-the-treatment-of-liver-cancer/299804

Surfactant-Based Anhydrous Nano Carrier System for Poorly Aqueous Soluble Anti-Cancer Drugs

Shekhar Verma, Nagendra Chandrawanshiand Vishal Jain (2021). *Handbook of Research on Advancements in Cancer Therapeutics* (pp. 413-432).

www.irma-international.org/chapter/surfactant-based-anhydrous-nano-carrier-system-for-poorly-aqueous-soluble-anti-cancer-drugs/267051

Predicting Dysfunctional Internet Use: The Role of Age, Conscientiousness, and Internet Literacy in Internet Addiction and Cyberbullying

Benjamin Stodt, Elisa Wegmannand Matthias Brand (2019). *Substance Abuse and Addiction: Breakthroughs in Research and Practice* (pp. 263-281).

www.irma-international.org/chapter/predicting-dysfunctional-internet-use/219419

Waiting for Health Care Services

Stefan Janzek-Hawlatand Hilda Tellolu (2017). *Transformative Healthcare Practice through Patient Engagement* (pp. 276-301).

www.irma-international.org/chapter/waiting-for-health-care-services/158995