

Chapter 44

Determination of Urban Growth by the Night–Time Images

Emre Yücer

Erzincan University, Turkey

Arzu Erener

Kocaeli University, Turkey

ABSTRACT

Urbanization in Turkey and in the world has increased in the last 30 years. Therefore, urban areas having a dynamic structure should be regularly monitored, and the urban region should be determined. There is a need of accurate current data that enables the monitoring of the temporally changing urban development and land use at regular intervals. In order to determine the urban sizes, the 2010 DMSP-OLS (US Air Force Meteorological Satellite Program – operational line scan system) nighttime images are used. This chapter is adapted to all provinces of Turkey. Nighttime images in the literature are mainly used for examining the global economic and demographic differences between countries. In order to determine the urban growth from nighttime images, two different image analyses including object-based and cell-based methods are used in this study. Object-oriented image analysis method includes image segmentation studies, and cell-based classification method contains natural breaks and OTSU method.

INTRODUCTION

Urbanization in Turkey and the world has continued to increase especially in the last 30 years. For developing countries, urbanization is a necessary phenomenon, especially in terms of the sustainability of urban growth. However, this growth must be controlled and in a planned manner, otherwise it could lead to many environmental problems. Therefore, urban areas, having a dynamic structure, should be monitored regularly and identified. Adequately accurate current data, which changes temporally and enables to monitor urban development and land use at regular intervals, are needed. Remote sensing technology is one of the most preferred methods in recent years to determine the urban areas developing fast and dynamically.

DOI: 10.4018/978-1-5225-7661-7.ch044

Determination of Urban Growth by the Night-Time Images

Within the scope of this study, using night-time images integrated with GIS, the spatial extent of urban areas in Turkey was identified and the methods used in the identification of the urban areas were compared. To determine the urban size of the cities in Turkey, DMSP- OLS (US Air Force Meteorological Satellite Program (Defense Meteorological Satellite Program) - Operational Line Scan System (Operational Linescan System) night-time images belonging to 2010 were used. While determining the urban size from night-time images, two different image analysis methods, object-based and pixel-based, were used. Object-based image analysis method involves image segmentation and pixel-based classification method involves natural breaks and Otsu method. By comparing the results obtained through these three methods, the method that gives the most accurate urban area size has been determined. Through the comparison of these methods' features used in the determination of the urban areas, the differences, advantages, and disadvantages of these methods have been revealed.

BACKGROUND

Urbanization is quickly increasing in the world and according to data of the United Nations Population Fund it is estimated that the world's population will be 12 billion by 2050, 80% of this population will be living in cities. In our country where an intensely urban migration from villages has been observed, there has been a clear increase in the urban areas. The effects of this increase occurring in urban areas on the social, economic and cultural characteristics of a city and population are inevitable.

Observation and evaluation of these changes occurring in urban areas are crucial for making balanced and sustainable urban development. Today, through the use of remote sensing and information technologies, the assessment process of data is accelerating and the productivity is increasing. In recent years, night-time images have been often used in the studies that determine the urban areas. The fact that these images can be accessed quickly and their assessment process is fast increases the interest in the night images.

The method was used to detect a threshold value for Chicago, Miami and Sacramento states in the USA. Maps of urban areas were created from night-time images in accord with this threshold value (Imhoff et al., 1997).

It has been showed that a single threshold value in night-time images has been more appropriate to determine the urban areas for different metropolitan regions. Therefore, Henderson et al. have indicated that the threshold value of the data in night-time images should be determined separately for each year for which a temporal analysis needs to be carried out for urban growth detection (Henderson et al., 2003).

As there is very little fluctuation in light intensity in Italy's Sicilian city at night, Elvidg et al. have implemented a quadratic regression model to inter-calibrate among years by using the data of the city as the reference data. The purpose of using a single city for calibration in night-time images is to correct the changes according to the years in the satellite and detector sensors (Elvidg et al., 2009).

By examining the global economic and demographic differences of night-time images, Levin and Duke conducted a study that could be used as an indication of the demographic and socio-economic characteristics of the residential area at a local scale. In the study, the population data of the settlements Israel and West Bank (Arab and Jewish) were compared. It has been stated that the economic and geo-political differences between Israel and the West Bank lead to pattern differences in night-time images (Levin & Duke, 2012).

11 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:
www.igi-global.com/chapter/determination-of-urban-growth-by-the-night-time-images/215894

Related Content

E-Participation and Citizen Relationship Management in Urban Governance: Tools and Methods

Jim P. Huebner (2015). *Public Affairs and Administration: Concepts, Methodologies, Tools, and Applications* (pp. 11-47).

www.irma-international.org/chapter/e-participation-and-citizen-relationship-management-in-urban-governance/127843

Participative Typologies: A Comparative Study Among Health Councils in Montevideo, Uruguay and Porto Alegre, Brazil

Andrea de Oliveira Gonçalves, Rodrigo de Souza Gonçalves and Elionor Farah Jreige Weffort (2011). *International Journal of Public and Private Healthcare Management and Economics* (pp. 40-52).

www.irma-international.org/article/participative-typologies-comparative-study-among/52797

Lighthouse in the Mist: State Governors' Crisis Communication on Twitter During the Early Stage of the COVID-19 Pandemic

Hanjin Mao, Meril Antony and Yujin J. Jung (2024). *International Journal of Public Administration in the Digital Age* (pp. 1-34).

www.irma-international.org/article/lighthouse-in-the-mist/344420

A Case Study on Implementing a Continuum of Care Based on Client Needs

Bonnie H.Y. Wong, William Y.W. Leung, Maria Chu and Kwong Y. Liu (2018). *International Journal of Public and Private Perspectives on Healthcare, Culture, and the Environment* (pp. 25-41).

www.irma-international.org/article/a-case-study-on-implementing-a-continuum-of-care-based-on-client-needs/207309

African Culture and Sustainability: The Case of the Grass Landers of Cameroon

Cornelius W. Wuchuan and Akoni Innocent Ngwainbi (2021). *International Journal of Public and Private Perspectives on Healthcare, Culture, and the Environment* (pp. 46-59).

www.irma-international.org/article/african-culture-and-sustainability/266293