

Chapter 1

Plasma Cocaine Metabolite Levels and Liver CYP450 3A4 Isoenzyme Activity as Indicators of Cocaine Metabolism in Rats Treated With Salako Supplements

Natwaine Sherune Gardner
University of the West Indies, Jamaica

Everard Barton
University of the West Indies, Jamaica

Kedon J. S. Luke
University of the West Indies, Jamaica

Lauriann E. Young-Martin
University of the West Indies, Jamaica

Andrew O. Wheatley
University of the West Indies, Jamaica

Ajibeke Salako-Akande
Getwele Natureceuticals, USA

Winston De La Haye
University of the West Indies, Jamaica

Henry Lowe
*Environmental Health Foundation,
Jamaica*

Perceval Steven Bahado-Singh
*University of Maryland Medical
System, USA*

Errol Morrison
*National Commission on Science and
Technology, Jamaica*

Lowell L. Dilworth
University of the West Indies, Jamaica

Denise Eldermire-Shearer
University of the West Indies, Jamaica

Donovan A. McGrowder
University of the West Indies, Jamaica

Helen Asemota
University of the West Indies, Jamaica

DOI: 10.4018/978-1-5225-5406-6.ch001

ABSTRACT

The effects of Salako nutritional supplements on cocaine-dependent Sprague Dawley rats was investigated. Rats were made cocaine-dependent using conditioned place preference (CPP) where craving was analyzed regularly. Cocaine metabolite levels were determined from blood samples. CYP450 3A4 isoenzyme activities were obtained using liver homogenate. Statistical analysis was done using SPSS one-way ANOVA and Duncans multiple range test. Results show that when cocaine use was discontinued, the supplements reduced craving of cocaine significantly. Blood plasma results showed higher benzoylecgonine equilibrium possibly indicating that the supplements aided the removal of stored cocaine metabolites which may have contributed to better management of craving in the rats. CYP450 3A4 isoenzyme activity was further enhanced by the supplements and is indicative of increased cocaine metabolism. The results indicate that the Salako nutritional supplements reduce craving caused by chronic cocaine administration by increasing the liver CYP450 3A4 isoenzyme activity, resulting in better plasma clearance.

INTRODUCTION

Cocaine (C₁₇H₂₁NO₄), classified as a stimulant, is described as being the most potent, powerfully addictive stimulant of natural origin. The leaves of the Coca plant (*Erythroxylum coca*) can be harvested several times a year. The alkaloid, cocaine, is extracted from the leaves of the Coca plant, which originates in South America, and to a lesser extent, in Africa, Indonesia and India (UNODC, 2010). Cocaine is one of the oldest known psychoactive substances. Coca use has been traced as far back as around 5000 B.C. wherein the leaves of the plant were continually chewed in the mouth. Pure cocaine was isolated in the 1880's (National Institute on Drug Abuse, 2008). In the early 1900s, pure cocaine was the main active ingredient in numerous pharmaceutical and recreational formulations due to their properties that enhanced general activity and decreased fatigue.

Illicit cocaine from South America is produced normally as relatively pure hydrochloride salt (ranging from 80 – 95 percent) for export to the United States. Illicit cocaine is distributed ranging from a white crystalline powder (cocaine hydrochloride), to that of an off-white chunky material ("crack" or "rock" cocaine) (Drug Enforcement Agency, 2005). Street cocaine is usually adulterated with various substances such as mannitol, lactose and glucose, talc and flour. Due to the heat

19 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/chapter/plasma-cocaine-metabolite-levels-and-liver-cyp450-3a4-isoenzyme-activity-as-indicators-of-cocaine-metabolism-in-rats-treated-with-salako-supplements/217760

Related Content

Heterogeneity Analysis in the University Context: A Proposal Based on Service Quality Perceptions

Maria Fuentes Blasco, Irene Gil Sauraand Beatriz Moliner Velásquez (2013). *Cases on Assessment and Evaluation in Education* (pp. 179-206).

www.irma-international.org/chapter/heterogeneity-analysis-university-context/69489

New Approaches of Nanocomposite Materials for Electromagnetic Sensors and Robotics

Alessandro Massaro, Fabrizio Spano, Diego Caratelli, Alexander Yarovoy, Roberto Cingolaniand Athanassia Athanassiou (2013). *Advanced Instrument Engineering: Measurement, Calibration, and Design* (pp. 57-73).

www.irma-international.org/chapter/new-approaches-nanocomposite-materials-electromagnetic/78170

AC Magnetic Measurements on Superconductors: Design of a Device for Magneto-Thermal Measurements

Philippe Laurent, Jean-François Fagnardand Philippe Vanderbemden (2011).

International Journal of Measurement Technologies and Instrumentation Engineering (pp. 53-66).

www.irma-international.org/article/magnetic-measurements-superconductors/62657

Efficient Artifact Elimination in Cardiac Signals using Variable Step Size Adaptive Noise Cancellers

S. Yasmin Fathima, G. V. S. Karthik, M. Zia Ur Rahmanand A. Lay-Ekuakille (2012).

International Journal of Measurement Technologies and Instrumentation Engineering (pp. 35-51).

www.irma-international.org/article/efficient-artifact-elimination-cardiac-signals/72700

Design of S-Band Antenna With L-Shaped Slits on Rectangular Patch With Defected Ground Structure

Ketavath Kumar Naik, Ravi Kumar Palla, Sriram Sandhya Raniand Dattatreya Gopi (2018). *International Journal of Measurement Technologies and Instrumentation Engineering* (pp. 50-58).

www.irma-international.org/article/design-of-s-band-antenna-with-l-shaped-slits-on-rectangular-patch-with-defected-ground-structure/217404