# Chapter 10 Tripartite and Quadpartite Size Ramsey Numbers for All Pairs of Connected Graphs on Four Vertices

Chula J. Jayawardene University of Colombo, Sri Lanka

### ABSTRACT

A popular area of graph theory is based on a paper written in 1930 by F. P. Ramsey titled "On a Problem on Formal Logic." A theorem which was proved in his paper triggered the study of modern Ramsey theory. However, his premature death at the young age of 26 hindered the development of this area of study at the initial stages. The balanced size multipartite Ramsey number mj (H,G) is defined as the smallest positive number s such that  $K_j \times s \rightarrow (H,G)$ . There are 36 pairs of (H, G), when H, G represent connected graphs on four vertices (as there are only 6 non-isomorphic connected graphs on four vertices). In this chapter, the authors find mj (H, G) exhaustively for all such pairs in the tripartite case j=3, and in the quadpartite case j=4, excluding the case  $m_4(K_4, K_4)$ . In this case, the only known result is that  $m_4$ ( $K_4, K_4$ ) is greater than or equal to 4, since no upper bound has been found as yet.

### HISTORY

After the publication of the original paper by F. P. Ramsey, the resurrection of Ramsey Theory for graphs emerged in the paper by Paul Erdös and George Szekeres, published around 1935.

Theorem (by Paul Erdös and George Szekeres)

For any  $m \ge 2$  and  $n \ge 2, r(m, n)$  exists and it satisfies

$$r(m,n) \le r(m-1,n) + r(m,n-1)$$

DOI: 10.4018/978-1-5225-9380-5.ch010

and

Following the publication of the paper by Paul Erdös and George Szekeres, the exact determination of these numbers for small graphs has been attempted by many mathematicians. Unfortunately, their progress was hindered by the stubbornly resistant (r(5,5)), which is presently known to lie between 43 and 48 (see Vigleik Angeltveit and Brendan D. McKay (Angelteveit, 2017) for the best upper bound of 48 and Exoo (Exoo, 1989) for the best lower bound of 43). Finding sharp bounds for (r(6,6)) appears to be an even more arduous task. This has been eloquently expressed by the great mathematician Paul Erdös using the following quotation (see 1990 Scientific American article by Ronald Graham and Joel Spencer). Suppose aliens invade the earth and threaten to obliterate it in a year's time unless human beings can find the Ramsey number for red five and blue five (r(5,5)). We could marshal the world's best minds and fastest computers, and within a year we could probably calculate the value. If the aliens demanded the Ramsey number for red six and blue six (r(6,6)), however, we would have no choice but to launch a pre-emptive attack. – Paul Erdös (1990 Scientific American article by Ronald Graham & Joel Spencer)

Researchers are now trying to approach this problem by using new techniques to investigate whether the lower bound 43 (Exoo, 1989) and the upper bound 48 (Angelteveit (2017)) could be improved. Off-shoots of classical Ramsey numbers, introduced by Burger and Vuuren (2004), Syfrizal et al., (2005, 2012), are the multipartite Ramsey numbers.

Let  $K_N$  denote the complete graph on N vertices. For any two graphs, say H, G (without loops and parallel edges), we say that  $K_N \to (H,G)$ , if for any red/blue colouring of  $K_N$ , given by  $K_N = H_R \oplus H_B$ , there exists a red copy H in  $H_R$  or a blue copy G in  $H_B$ . In Ramsey theory one important part deals with the exact determination of Ramsey numbers, r(H,G), defined as the smallest positive number N such that  $K_N \to (H,G)$ . Formally, the balance multipartite graph  $K_{j\times s}$  can be viewed as a graph consisting of the vertex set

$$V\left(K_{_{j\times s}}\right) = \left\{v_{_{m,n}}: m \in \left\{1,2,\ldots j\right\} and \ n \in \left\{1,2,\ldots s\right\}\right\}$$

and the edge set

$$E\left(K_{_{j\times s}}\right) = \left\{ \left(v_{_{m,n}}, v_{_{m',n'}}\right) : m, m' \in \left\{1, 2, \dots, j\right\}, m \neq m \text{ and } n, n' \in \left\{1, 2, \dots, s\right\} \right\}.$$

In such a graph, the set of vertices in the  $m^{th}$  partite set is denoted by  $\{v_{mn} \mid n \in \{1, 2, ..., s\}\}$ . The Ramsey number  $m_i(H,G)$  is defined as the smallest positive number s such that  $K_{ixs} \rightarrow (H,G)$ .

14 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/tripartite-and-quadpartite-size-ramsey-numbers-

for-all-pairs-of-connected-graphs-on-four-vertices/235539

### **Related Content**

## On the Influence of Fractional Derivative on Chaos Control of a New Fractional-Order Hyperchaotic System

Ahmed Ezzat Mohamed Matouk (2020). Advanced Applications of Fractional Differential Operators to Science and Technology (pp. 115-132).

www.irma-international.org/chapter/on-the-influence-of-fractional-derivative-on-chaos-control-of-a-new-fractional-orderhyperchaotic-system/255469

### **Digital Beauty**

(2024). Utilizing Visuals and Information Technology in Mathematics Classrooms (pp. 218-236). www.irma-international.org/chapter/digital-beauty/346428

### Review of Basic Material: Induction and Counting

(2025). *Mathematics for Effective Management (pp. 1-24).* www.irma-international.org/chapter/review-of-basic-material/368862

#### Joint Normal Partitions and Hierarchical Filling of N-Dimensional Spaces

(2021). Normal Partitions and Hierarchical Fillings of N-Dimensional Spaces (pp. 261-272). www.irma-international.org/chapter/joint-normal-partitions-and-hierarchical-filling-of-n-dimensional-spaces/267849

### On NeutroQuadrupleRings

Muritala Abiodun Ibrahim, Agboola Adesina Abdul Akeem, Zulaihat Hassan-Ibrahimand Emmanuel O. Adeleke (2022). *Theory and Applications of NeutroAlgebras as Generalizations of Classical Algebras (pp. 29-57).* 

www.irma-international.org/chapter/on-neutroquadruplerings/302850