Chapter 7 Fundamental Theories and Kinetic Models for the Pyrolysis of Lignocellulosic Biomass Wastes

Olagoke Oladokun

University of Maiduguri, Nigeria

Bemgba B. Nyakuma

Universiti Teknologi Malaysia, Malaysia

Arshad Ahmad

Universiti Teknologi Malaysia, Malaysia

ABSTRACT

Biomass agricultural waste has a great potential for meeting part of the world energy need and is completely environmentally friendly. One conversion method is thermochemical processes and specifically, pyrolysis. Pyrolysis converts the lignocellulose waste to fuel and essential chemicals into three products: biogas, bio-oil, and biochar. However, performance issues limit the potential of lignocellulose pyrolysis such as design and operation of pyrolysis reactor for effective heat transfer from the heat source to the biomass feedstock. Therefore, this study presents the necessary tools for pyrolysis scientists and engineers in determining the optimal operation and design of lignocellulose agricultural waste pyrolysis. The tools consist of mathematical equations that govern the lignocellulose kinetics (model and model-free) and pyrolysis reactor macro and micro models. A practical model for hydrogen production from pyrolysis bio oil solidifies the viability of biomass as an energy source.

DOI: 10.4018/978-1-7998-0369-0.ch007

INTRODUCTION

Biomass is an abundant, renewable and sustainable source of energy for the future. Biomass typically includes agricultural waste residues, forest waste products, municipal solid waste and energy crops such as perennial grasses (Hayes, 2013; Liew, Hassim, & Ng, 2014; Mohammadi et al., 2011). Biomass has been burnt directly to generate heat for household, gasified to power engines or converted to biofuels and syngas by pyrolysis and other thermochemical processes (Ahmed et al., 2015; Basu, 2013; A. V. Bridgwater, 2012; Miller & Bellan, 1997). A vast amount of biomass is produced every year and of that only 5% (13.5 billion metric tonnes) are effectively converted to energy, which is equivalent to about 26% of the world's energy consumption (Basu, 2013). Figure 1 gives the biomass resources and their energy contribution (Balat, 2008; Demirbas, 2007; L. Zhang, Xu, & Champagne, 2010)).

However, the conversion of agroforest and agricultural product into bioenergy and biofuels presents significant challenges for the environment and human food supply (Oladokun et al., 2017; Sims, Mabee, Saddler, & Taylor, 2010). Consequently, the search for alternative biomass feedstocks such as perennial energy grasses must continue. Numerous studies have investigated the carbonization, torrefaction, gasification and pyrolysis potential of energy grasses such Miscanthus, Switchgrass, and Canary Reed (Floudas, Elia, & Baliban, 2012; Heaton et al., 2010; Lemus et al., 2002; Moutsoglou, 2012; Woli et al., 2011; Yang, Yan, Chen, Lee, & Zheng, 2007). Other potential energy grasses such Elephant grass, and *Imperata cylindrica* found around the world can also be utilised for bioenergy applications.

The conversion of biomass resources into solid, liquid and gaseous products such as biohydrogen through biochemical or thermochemical routes is widely reported in the literature (Monlau et al., 2013; Nyakuma et al., 2014; X. Zhang, Wang, Ma, Zhang, & Jiang, 2013). The biochemical route involves the breakdown of biomass molecules by enzymes or acid (enzymatic or acid hydrolysis) into sugar and subsequent fermentation to bioethanol (Balan et al., 2012; Basu, 2013). The process is well established and cost effective due to advances in research and optimisation of fermentation. However, biochemical processing of biomass into fuels cannot be easily integrated into existing commercial refining operation compared to thermochemical routes (Anthony V. Bridgwater, 2012). Combustion, pyrolysis and gasifi-




Figure 1. Biomass resources and energy contribution

27 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/fundamental-theories-and-kinetic-models-for-the-pyrolysis-of-lignocellulosic-biomass-wastes/242014

Related Content

Genetically Engineered Microorganisms for Bioremediation Processes: GEMs for Bioremediaton

Stephen Rathinaraj Benjamin, Fabio de Limaand Ashok K. Rathoure (2016). *Toxicity and Waste Management Using Bioremediation (pp. 113-140).*

www.irma-international.org/chapter/genetically-engineered-microorganisms-for-bioremediation-processes/141796

Implications of Molecular Docking Assay for Bioremediation

Zarrin Basharat, Monazza Bibiand Azra Yasmin (2017). Handbook of Research on Inventive Bioremediation Techniques (pp. 24-45).

www.irma-international.org/chapter/implications-of-molecular-docking-assay-for-bioremediation/176457

Influence of Joint Orientation on the Behavior of Dam Foundation Resting on Jointed Rock Mass Under Earthquake Loading Condition

Prasun Halder, Saurabh Kumar, Bappaditya Mannaand K. G. Sharma (2019). *International Journal of Geotechnical Earthquake Engineering (pp. 1-16).*

www.irma-international.org/article/influence-of-joint-orientation-on-the-behavior-of-dam-foundation-resting-on-jointed-rock-mass-under-earthquake-loading-condition/225086

Landslides: A Guide to Researching Landslide Phenomena and Processes

Snježana Mihali Arbanasand Željko Arbanas (2015). *Handbook of Research on Advancements in Environmental Engineering (pp. 474-510).*

www.irma-international.org/chapter/landslides/122644

Estimation of Peak Ground Acceleration and Its Uncertainty for Northern Indian Region

Girish Chandra Joshiand Mukat Lal Sharma (2011). *International Journal of Geotechnical Earthquake Engineering (pp. 1-19).*

www.irma-international.org/article/estimation-peak-ground-acceleration-its/52783