# Chapter 7 On the Orthogonality of the q-Derivatives of the Discrete q-Hermite I Polynomials

Sakina Alwhishi El Mergib University, Libya

**Rezan Sevinik Adıgüzel** *Atılım University, Turkey* 

Mehmet Turan Atilim University, Turkey

#### ABSTRACT

Discrete q-Hermite I polynomials are a member of the q-polynomials of the Hahn class. They are the polynomial solutions of a second order difference equation of hypergeometric type. These polynomials are one of the q-analogous of the Hermite polynomials. It is well known that the q-Hermite I polynomials approach the Hermite polynomials as q tends to 1. In this chapter, the orthogonality property of the discrete q-Hermite I polynomials is considered. Moreover, the orthogonality relation for the k-th order q-derivatives of the discrete q-Hermite I polynomials is obtained. Finally, it is shown that, under a suitable transformation, these relations give the corresponding relations for the Hermite polynomials in the limiting case as q goes to 1.

DOI: 10.4018/978-1-7998-0134-4.ch007

Copyright © 2020, IGI Global. Copying or distributing in print or electronic forms without written permission of IGI Global is prohibited.

#### INTRODUCTION

The so-called *q*.polynomials have great applications in several problems on theoretical and mathematical physics, for example, in the continued fractions, Eulerian series (Fine, 1988), algebras and quantum groups (Koornwinder, 1995, 1994; Vilenkin & Klimyk, 1998), discrete mathematics, algebraic combinatorics (coding theory, design theory, various theories of group representation) (Bannai, 1990), *q*.Schrödinger equation and *q*.harmonic oscillators (Askey et al., 1993; Askey & Suslov, 1993; Atakishiyev & Suslov, 1991; Berg & Ismail, 1994; Borzov & Damaskinsky, 2003; Macfarlane, 1989).

Discrete q. Hermite I polynomials form an important class of q. polynomials in the Hahn sense. They satisfy the following q. difference equation of hypergeometric type:

$$-D_{q}D_{q^{-1}}y + \frac{x}{1-q}D_{q^{-1}}y + \lambda y = 0,$$
(1)

where  $\lambda$  is a constant which has a particular form depending on the parameter q. and  $D_q$  is the q.Jackson derivative (Koekoek et al., 2010; Nikiforov et al., 1991; Nikiforov & Uvarov, 1986, 1988).

In fact, discrete *q*.Hermite polynomials are the *q*.analogous of the Hermite polynomials which are one of the important orthogonal family of the classical orthogonal polynomials (Koekoek et al., 2010; Nikiforov et al., 1991; Nikiforov & Uvarov, 1986, 1988; Szegö, 1939) that have enormous applications in mathematics and physics. They satisfy the following differential equation of hypergeometric type

$$y'' - 2zy' + 2ny = 0, (2)$$

where  $n \in \mathbb{N}_0$ . The Hermite polynomials and their *q*. analogues can be obtained by using an appropriate limit relation from the other orthogonal polynomials (Andrews et al., 1999; Koekoek et al., 2010).

The classical orthogonal polynomials and their *q*.analogues have several properties. For example the differential and the difference equations that they satisfy have special forms, which are studied by Bochner in 1929 (Bochner, 1929) and by Routh in 1885 (Routh, 1885). Rodrigues formula is another property that provides a representation for the classical polynomials derived by Tricomi (Tricomi, 1955) and (Cryer, 1970). The three terms recurrence relation (TRRR) is also a way to construct classical orthogonal polynomials considered by Chihara (Chihara, 1978) and Szegö (Szegö, 1939). Moreover, the generating function is another option to define these

26 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igiglobal.com/chapter/on-the-orthogonality-of-the-q-derivativesof-the-discrete-q-hermite-i-polynomials/242345

#### **Related Content**

## Bio-Inspired Techniques in the Clustering of Texts: Synthesis and Comparative Study

Reda Mohamed Hamou, Hadj Ahmed Bouararaand Abdelmalek Amine (2015). International Journal of Applied Metaheuristic Computing (pp. 39-68). www.irma-international.org/article/bio-inspired-techniques-in-the-clustering-of-texts/132523

## New Evolutionary Algorithm Based on 2-Opt Local Search to Solve the Vehicle Routing Problem with Private Fleet and Common Carrier

Jalel Euchi, Habib Chabchouband Adnan Yassine (2011). *International Journal of Applied Metaheuristic Computing (pp. 58-82).* www.irma-international.org/article/new-evolutionary-algorithm-based-opt/52793

## (Approximate) Frequent Item Set Mining Made Simple with a Split and Merge Algorithm

Christian Borgeltand Xiaomeng Wang (2010). *Scalable Fuzzy Algorithms for Data Management and Analysis: Methods and Design (pp. 254-272).* www.irma-international.org/chapter/approximate-frequent-item-set-mining/38572

### Development of Energy Efficient and Optimized Coverage Area Network Configuration to Achieve Reliable WSN Network Using Meta-Heuristic Approaches

Avishek Banerjee, Victor Das, Arindam Biswas, Samiran Chattopadhyayand Utpal Biswas (2021). *International Journal of Applied Metaheuristic Computing (pp. 1-27).* www.irma-international.org/article/development-of-energy-efficient-and-optimized-coveragearea-network-configuration-to-achieve-reliable-wsn-network-using-meta-heuristicapproaches/284417

# A Formal Approach to Evaluating Medical Ontology Systems using Naturalness

Yoo Jung An, Kuo-Chuan Huang, Soon Ae Chunand James Geller (2012). Innovations in Data Methodologies and Computational Algorithms for Medical Applications (pp. 1-18).

www.irma-international.org/chapter/formal-approach-evaluating-medical-ontology/65148