


Chapter 10

Antioxidative Agents From Medicinal Plants

Sandeep Kumar

 <https://orcid.org/0000-0002-7576-9696>

National Dairy Research Institute, India

Ahmad Hussain

 <https://orcid.org/0000-0002-1603-0154>

National Dairy Research Institute, India

Manish Singh Sansi

National Dairy Research Institute, India

Daraksha Iram

National Dairy Research Institute, India

Priyanka

Indian Institute of Technology, Delhi, India

Ashutosh Vats

National Dairy Research Institute, India

ABSTRACT

The medicinal plants have been used by humans since ancient times, and the great civilizations of the world in ancient times were well aware of the benefits brought by the use of medicinal plants. This chapter provides important information regarding medicinal plants that have a wide variety of antioxidative agents ranging from bitter compounds that stimulate digestion system, phenolic compounds for antioxidant and numerous other pharmacological properties, antibacterial, and antifungal to tannins that act as natural antibiotics, diuretic substances, alkaloids, and so forth.

DOI: 10.4018/978-1-6684-3546-5.ch010

INTRODUCTION

Medicinal plants are utilized by humans since history and also the great civilizations of the world were well aware of their use. However, the use of medicinal plants keeps on increasing after understanding their detailed functions and the components present in them. The use of medicinal plants were used by adding plant parts or extracts as additives in different types of foods and various herbal drinks for enhancing their effectiveness. Antioxidants are the substances which act against oxidative reaction. Within the biological systems, the antioxidant is the biological active molecule that is capable of preventing a precise oxidation reaction catalyzed by enzymes or a material which reacts with oxidizing agents before it causes oxidative damage to the other molecules.

Antioxidants even in trace amounts can stop or pause the oxidation process of easily oxidizable biomolecules present in the cell (Brewer, 2011). Free radicals are generated in the form of reactive oxygen species within the living organisms by completely different biochemical and physiological pathways has been observed, however, increased production of those may result in toward the progress of various diseases (Halliwell, 2007). Reactive oxygen species generation is necessary for many physiological processes e.g. cell proliferation, differentiation and programmed pathway for death of cell. They are intermediates of many important signaling processes; however, they are also toxic byproducts of normal metabolism (Mittler, 2017).

Reactive oxygen species are generated by the incomplete reduction of environmental oxygen (O_2) i.e., by excitation of oxygen to reactive half singlet oxygen (O_2^1) or by the transfer of electrons to oxygen which in turn forms superoxide radical ($O_2^{\cdot-}$), hydroxy radical (HO^{\cdot}) or hydrogen peroxide (H_2O_2). Reactive oxygen species can cause oxidation of cellular organelles and results in the destruction of cells and tissues by oxidative processes (Asada & Takahashi 1987).

We have identified many potential sources of ROS generation in plants. Some of them are the normal reactions going down within the cell and concerned in normal metabolisms, such as respiration and photosynthesis. Reactive oxygen species may also be created by alternative factors like pathways increased throughout abiotic stress e.g. glycolate oxidase enzyme in peroxisomes during the process of photorespiration. Few new sources of reactive oxygen species generation have been identified in plants e.g. amine oxidases, peroxidases bounded to cell wall, and NADPH oxidases. These processes are tightly regulated and participates in reactive oxygen species generation during apoptosis and defense from pathogen (Asada & Takahashi, 1987).

The medicinal plants are simply accessible and great supply of antioxidative compound as they contains a blend of various chemical substances that can demonstrate individually or unitedly to fix health problems and improvement in diseases. Indeed, a solitary plant might have a decent variety of phytochemical compounds starting from phenolic compounds as antioxidant and bitter compounds which stimulates digestion, also and various alternative medicinal properties, antifungal, antibacterial, diuretic substances tannins that act as naturally occurring antibiotics, and, alkaloids, and so forth (Miguel, 2010). In the body, antioxidative agents decrease the hazard of degenerative ailments emerging due to oxidative pressure. The antioxidative agents can be of different types based on sources i.e. primary antioxidative agents and secondary antioxidative agents present in the nature. The primary antioxidants are those that counterbalance reactive oxygen species through a single electron transfer system or by either donating a hydrogen particle. On the opposite hand, secondary antioxidants are biological compounds that have the ability to neutralize pro-oxidant catalysts. These include chelators which bind tightly to pro-oxidant metallic ions such as copper and iron, demonstrated by citric acid (CA) and ethylenediaminetetraacetic

14 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/antioxidative-agents-from-medicinal-plants/289481

Related Content

A Strategic Perspective on Using Symbolic Transformation in STEM Education: Robotics and Automation

Jack M. Rappaport, Stephen B. Richter and Dennis T. Kennedy (2018). *Applications of Neuroscience: Breakthroughs in Research and Practice* (pp. 242-284).

www.irma-international.org/chapter/a-strategic-perspective-on-using-symbolic-transformation-in-stem-education/199639

Nanoparticle-Mediated Therapeutic Approach for Ulcerative Colitis Treatment

Subhamoy Banerjee (2021). *Diagnostic and Treatment Methods for Ulcerative Colitis and Colitis-Associated Cancer* (pp. 96-117).

www.irma-international.org/chapter/nanoparticle-mediated-therapeutic-approach-for-ulcerative-colitis-treatment/274081

Visual Communication to Improve Relationship Quality in Spousal Caregivers of Individuals With Alzheimer's Disease

Nola Freeman (2019). *Chronic Illness and Long-Term Care: Breakthroughs in Research and Practice* (pp. 59-76).

www.irma-international.org/chapter/visual-communication-to-improve-relationship-quality-in-spousal-caregivers-of-individuals-with-alzheimers-disease/213337

Chronic Condition Management Using Remote Monitoring and Telehomecare

Maximillian E. Stachura, Ellen B. Bolch, Katherine F. Piette and Joseph F. Ebberwein (2019). *Chronic Illness and Long-Term Care: Breakthroughs in Research and Practice* (pp. 669-686).

www.irma-international.org/chapter/chronic-condition-management-using-remote-monitoring-and-telehomecare/213375

Contemporary Heart Failure Treatment Based on Improved Knowledge and Personalized Care of Comorbidities

Kostas Giokas, Charalampos Tsirmpas, Athanasios Anastasiou, Dimitra Iliopoulou, Vassilia Costarides and Dimitris Koutsouris (2016). *Handbook of Research on Trends in the Diagnosis and Treatment of Chronic Conditions* (pp. 301-314).

www.irma-international.org/chapter/contemporary-heart-failure-treatment-based-on-improved-knowledge-and-personalized-care-of-comorbidities/136523