

Chapter 1

Clinical Decision Support System to Prevent Toxicity in Patients Treated With Digoxin

Asunción Albert

University Hospital Dr. Peset/University of Valencia, Spain

Antonio J. Serrano

University of Valencia, Spain

Emilio Soria

University of Valencia, Spain

Nicolás Victor Jiménez

University Hospital Dr. Peset/University of Valencia, Spain

ABSTRACT

Digoxin (DGX) is drug used to control signs and symptoms involved in congestive heart failure and atrial fibrillation. Due to its narrow therapeutic range more than 10% of the patients treated with DGX can suffer toxic effects, but it is estimated that half of the cases of digitalis toxicity could be prevented. Two multivariate models were developed to prevent digitalis toxicity, with data of 125 patients monitored at the Pharmacy Service of the University Hospital Dr. Peset (Valencia, Spain). One hundred and six patients were used to develop the models and the other (55) to validate them. The logistic regression model achieved a 90.9% of sensitivity and 81.8% of specificity in the validation set with only 2 variables that are statistical significant (cardiomegaly and digoxin plasmatic concentration). The feed-forward multilayer network model achieved 100% of sensitivity and 90.9% of specificity equal in the validation set but all 14 variables studied are used as input in this model. A software for identifying patients at risk in order to implement preventive measures to avoid sings and symptoms of digitalis toxicity was developed.

DOI: 10.4018/978-1-61520-977-4.ch001

INTRODUCTION

Digoxin is one of the most widely prescribed drugs for the treatment of congestive heart failure and atrial arrhythmia. Despite current guidelines relegate digoxin use to patients that remain symptomatic after receiving optimum doses of antihypertensive and the utility of digoxin in atrial dysrhythmia is only superior to other agents when the patient has associated heart failure.

Digoxin use has decreased significantly from 31.4% (late 2001) to 23.5% in (late 2004) ($p < 0.00001$), the number of toxic or potentially toxic exposures to digoxin requiring hospitalization has not decreased (Hussain, Swindle, & Hauptman, 2006), specially in geriatric patients who uses about 80% of the global consumption of digitalis drugs (Kernan, Castellsague, Perlman, & Ostfeld, 1994; Warren, McBean, Hass, & Babish, 1994).

The Institute for Safe Medication Practices classifies digoxin as a “high alert medication”; that is drugs that bear a heightened risk of causing significant patient harm when they are used in error (*High-alert medication list*).

Therefore, digoxin intoxication represents about a fourth of the adverse drug events (ADEs) detected in the geriatric population, which frequently results in hospitalization (Gurwitz et al., 2000). Digoxin is the fourth drug class (7.7%) most frequently involved in potentially preventable adverse drug events at the emergency room (Otero et al., 2006), and one of the ten drugs most frequently associated with medication that demand interdisciplinary (physicians, pharmacists and nurses) actions (Llopis, Albert, Sancho, Calipienso, & Jiménez-Torres, 1999; Markota, Markota, Tomic, & Zelenika, 2009; Mutnick et al., 1997; Rupawala, Kshirsagar, & Gogtay, 2009).

The wide pharmacokinetic and pharmacodynamic variability of digoxin due to its narrow therapeutic range, together with the multiple factors that can affect the organism response to digoxin and the diversity of clinical manifestations

of digitalis toxicity (DT), results in digoxin being frequently associated to drug related problems. It is estimated that half of the cases of DT could have been prevented; this gives an opportunity to improve the treatment that patients receive and reduce medical costs (Gandhi, Vlasses, Morton, & Bauman, 1997).

In order to improve efficiency, with the available resources, it is necessary to identify those patients that are most susceptible to profit from pharmaceutical care actions; that is to say, to identify patients with either potential or real drug related problems, in order to prevent the former and resolve the latter. Multivariate statistical methods, as logistic regression and neural networks, are useful tools to develop prediction models of morbidity (toxicity or therapeutic failure) associated to drug use, particularly when factors that influence the therapeutic response are analyzed.

Neural networks technology does not require “a priori” models and represents an important improvement to the multivariate analysis. Artificial neural networks stand out for being able to find complex relations between dependent variables and the independent ones, in agreement with their superior ability to discover non linear relations, a representative characteristic of pharmacologic variability. The application of these models presents advantages over the actual statistical methods, although this doesn't mean not to recognize their limitations (Soria, Jiménez, & Serrano, 2000).

Neural Networks in Clinical Medicine

Artificial neural networks have been applied with marked success in classification problems, modeling, signal processing and time series prediction in as scattered fields as engineering, economics and telecommunications. Experts in other fields have recognized the new technology potential and they are incorporating it into their work. However, in health sciences, statistical methods are still mainly used (Michie, Spiegelhalter, & Taylor, 1994), such as logistic regression applied to data analysis. In

19 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:
www.igi-global.com/chapter/clinical-decision-support-system-prevent/43247

Related Content

E-Health Dot-Coms' Critical Success Factors

Abrams A. O'Buyonge (2009). *Medical Informatics: Concepts, Methodologies, Tools, and Applications* (pp. 1813-1821).

www.irma-international.org/chapter/health-dot-coms-critical-success/26338

Sensing of Vital Signs and Transmission Using Wireless Networks

Yousef Jasebian (2009). *Mobile Health Solutions for Biomedical Applications* (pp. 180-207).

www.irma-international.org/chapter/sensing-vital-signs-transmission-using/26772

Drowsiness Detection by the Systems Dynamic Approach of Oculomotor Systems

Dabhu Suman, Malini Mudigonda, B. Ram Reddy and Yashwanth Vyza (2022). *International Journal of Biomedical and Clinical Engineering* (pp. 1-27).

www.irma-international.org/article/drowsiness-detection-by-the-systems-dynamic-approach-of-oculomotor-systems/295866

Case Based Reasoning for Customizing Treatment Processes

Carolyn Kaiser (2009). *Handbook of Research on Distributed Medical Informatics and E-Health* (pp. 351-366).

www.irma-international.org/chapter/case-based-reasoning-customizing-treatment/19945

EEG Synchronization and Brain Networks: A Case Study in Fatigue

Anwasha Sengupta, Subhadeep Datta, Sibsambhu Karand Aurobinda Routray (2015). *International Journal of Biomedical and Clinical Engineering* (pp. 1-11).

www.irma-international.org/article/eeg-synchronization-and-brain-networks/138223