

Chapter 5

Mass and Heat Transfer Modeling

Ronald W. Breault

National Energy Technology Laboratory, USA

ABSTRACT

Mass and heat transfer are important to reactor modeling using CFD. Reactants (mass) move within the flow structure in order to react with other species and thereby form the products, both desirable and undesirable. This movement in mass occurs either by diffusion or by turbulent dispersion. In a similar fashion, heat is transferred from one point in the flow structure to another point by convective transfer between the phases and with the translational effect that occurs with the turbulent dispersion of the mass. In addition, heat can be transferred across the system boundary. In all these cases, fundamental mechanistic models are put forward that can be incorporated in the CFD code to calculate these transfer properties based upon the local hydrodynamic conditions. The chapter is organized such that dense phase systems are covered first and then dilute phase systems. Within each of these areas mass transfer is covered first and followed by heat transfer. The topics are covered in the following order: Diffusional mass transfer, Turbulent dispersion, Convective heat transfer between phases and Convective heat transfer at the system boundary.

INTRODUCTION

Prior to jumping into the discussion of the diffusional mass transfer, heat transfer and the turbulent dispersion, a short overview of the balances contained within MFIX is shown below such that these properties can be identified. Interphase mass transfer occurs by two mechanisms in heterogeneous gas-solids flows. The first of these is through diffusion across the boundary layer and the other is from turbulent transport of the species in gas eddies or solids clusters. Before to getting into the details of these two mechanisms

DOI: 10.4018/978-1-61520-651-3.ch005

and the methods which will be used to calculate these quantities, it is important to understand how and where these properties are used in reacting computational multiphase flow fluid dynamics.

For isothermal conditions, the continuity, momentum, and species balance equations are given below.

Gas-Phase Continuity

$$\frac{\partial}{\partial t}(\epsilon_g \rho_g) + \nabla \circ (\epsilon_g \rho_g \vec{v}_g) = \sum_{n=1}^{N_g} R_{gn} \quad (1)$$

Solids-Phase Continuity

$$\frac{\partial}{\partial t}(\epsilon_s \rho_s) + \nabla \circ (\epsilon_s \rho_s \vec{v}_s) = \sum_{n=1}^{N_s} R_{sn} \quad (2)$$

Gas-Phase Momentum

$$\frac{\partial}{\partial t}(\epsilon_g \rho_g \vec{v}_g) + \nabla \circ (\epsilon_g \rho_g \vec{v}_g \vec{v}_g) = -\epsilon_g \nabla P_g + \nabla \circ \overline{\tau}_g + F_{gs}(\vec{v}_s - \vec{v}_g) + \epsilon_g \rho_g \vec{g} - R_0(\xi_0 \vec{v}_g + \overline{\xi_0} \vec{v}_s) \quad (3)$$

Solids-Phase Momentum

$$\frac{\partial}{\partial t}(\epsilon_s \rho_s \vec{v}_s) + \nabla \circ (\epsilon_s \rho_s \vec{v}_s \vec{v}_s) = -\epsilon_s \nabla P_g + \nabla \circ \overline{S}_m - F_{gs}(\vec{v}_s - \vec{v}_g) + \epsilon_s \rho_s \vec{g} - R_0(\xi_0 \vec{v}_g + \overline{\xi_0} \vec{v}_s) \quad (4)$$

Species Balance

$$\frac{\partial}{\partial t}(\epsilon_m \rho_m X_{mn}) + \nabla \circ (\epsilon_m \rho_m X_{mn} \vec{v}_m) = \nabla \circ D_{mn} \nabla X_{mn} + R_{mn} \quad (5)$$

where $m = g$ or s for the gas or solids phase, D_{mn} is the turbulent diffusive term, R_{mn} is the reaction rate which is dependent upon the diffusional mass transfer, and $\overline{\xi_0} = 1 - \xi_0$, $\xi_0 = 1$ if $R_0 < 0$ else $\xi_0 = 0$. The eight dependent hydrodynamic variables in three dimensions are void fraction ϵ_g (the solids fraction $\epsilon_s = 1 - \epsilon_g$), pressure P_g , and six velocity components. These are found by using MFIX to numerically solve the coupled non-linear partial differential equations. The number of species mass fractions (X_{mn}) tracked are given in the chemistry model. Constitutive relations needed to close the system, and the gas/solids energy balance equations can be found in the following references (Syamlal, M., Rogers, W., & O'Brien, T. (1993); Syamlal, M.(1998)). A discussion on the solution procedure and further numerical references was prepared by Guenther & Syamlal (2001).

In general, the reaction rate or kinetics follow a conventional resistance model pathway, where the rate is proportional to the sum of the resistance elements such as transfer from the bulk to the particle surface plus the internal resistance which may include diffusion of the gas reactant in through an ash/

23 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/mass-heat-transfer-modeling/47493

Related Content

Graphene-Based Sensors for Monitoring Strain: A First-Principles Density Functional Theory Analysis

M. Mirnezhad, R. Ansari, H. Rouhiand M. Faghihnasiri (2013). *International Journal of Chemoinformatics and Chemical Engineering* (pp. 74-83).

www.irma-international.org/article/graphene-based-sensors-monitoring-strain/75224

Laminar Mixed Convection in Two Concentric Vertical Cylinders

Ahmed Hadjadjand Seghir Maamir (2016). *International Journal of Chemoinformatics and Chemical Engineering* (pp. 24-43).

www.irma-international.org/article/laminar-mixed-convection-in-two-concentric-vertical-cylinders/166582

Experimental Research on Grinding Temperature With Different Workpiece Materials

(2020). *Enhanced Heat Transfer Mechanism of Nanofluid MQL Cooling Grinding* (pp. 226-248).

www.irma-international.org/chapter/experimental-research-on-grinding-temperature-with-different-workpiece-materials/247317

Preparation of a Uranium Conversion Plant's Nuclear Waste for Final Disposal by Means of Magnetically Assisted Chemical Separation

Ahad Ghaemi, Mehdi Maghsudi, Fatemeh Hanifpourand Mohammad Samadfam (2013). *International Journal of Chemoinformatics and Chemical Engineering* (pp. 8-18).

www.irma-international.org/article/preparation-uranium-conversion-plant-nuclear/75220

Normal and Deteriorated Heat Transfer Under Heating Turbulent Supercritical Pressure Coolants Flows in Round Tubes

Vladimir A. Kurganov, Yury A. Zeigarnikand Irina V. Maslakova (2021). *Handbook of Research on Advancements in Supercritical Fluids Applications for Sustainable Energy Systems* (pp. 494-532).

www.irma-international.org/chapter/normal-and-deteriorated-heat-transfer-under-heating-turbulent-supercritical-pressure-coolants-flows-in-round-tubes/259848