

Chapter 3.6

Confirmatory Factor Analysis to Establish Determinants of Wireless Technology in the Indian Healthcare

Raj Gururajan

University of Southern Queensland, Australia

Tiana Gurney

University of Southern Queensland, Australia

Abdul Hafeez-Baig

University of Southern Queensland, Australia

ABSTRACT

This study reports the determinants of wireless technology in the Indian healthcare validated by a second order regression model. In order to assert the determinants, a qualitative study was conducted with 30 physicians using interviews to arrive at a set of barriers and drivers. Further analysis of the qualitative data indicated that there is a third component emerging, namely, clinical influence. The interview data was used to develop a survey instrument and this was administered on the Indian clinicians with 200 completed surveys. This data was used to establish the sub-components of the three major determinants as identified in the qualitative study. This is reported in this article.

INTRODUCTION

In the last few years, high expectations, technological developments, and effective and efficient services have been shown to be prerequisites for improvements in the healthcare domain (Rogoski, 2005; Versel, 2008). Latest trends in the healthcare sector include the design of more flexible and efficient service provider frameworks aimed at providing health services to all stakeholders. In order to implement such frameworks, wireless technology is increasingly being used in the healthcare sector. A decrease in the cost of wireless devices and improved awareness of the benefits by using related wireless applications are two of the contributing factors towards the increased use of wireless technology in this sector (Gururajan,

Quaddus, Fink, Vuori, & Soar, 2005; R. Gururajan, Hafeez-Baig, & Gururjan, 2008). Even though the future of this technology and its usability is promising, its adoption is still in its infancy, which is attributed to the complex and critical nature of the healthcare environment. In the current competitive and complex business environment, technology developments have played a critical role in delivering high quality of care (Reinecke, 2004). However, there is limited knowledge and empirical research on the effectiveness and adoption of wireless technology in general, and in the Indian healthcare system in particular.

Recent research has established that investment in emerging Information Technology (IT), including Information Systems (IS), can lead to productivity gains only if they are accepted and effectively used by respective stakeholders. Consequently, acceptance and utilization of IT/IS in the healthcare environment have been central themes in the information systems literature. Therefore, the fundamental focus of this research is to investigate and examine the influence of internal and external determinants on the usefulness of wireless technology. Further, this research also assesses how its acceptance contributes to the adoption of wireless technology. We believe that this research is the first of its kind attempted in the Indian healthcare domain and it employs empirical evidence to explore the impact of wireless technology and its usefulness in the Indian healthcare system. The Indian healthcare domain is at the forefront in adopting the latest medical technologies and applications, as evidenced by media reports and, as such, it constitutes an excellent context for validating existing adoption theories and extending them.

The main contribution of this research includes the identification of a set of drivers and barriers to using wireless technology in a given Indian healthcare setting. In addition to this, for the first time, a set of clinical factors influencing the adoption of wireless technology has been identified and validated using a second order regression model.

LITERATURE REVIEW

The concept of wireless technology in healthcare is discussed in many studies (Dyer, 2003; Hu, Chau, & Sheng, 2002; Sausser, 2003; Simpson, 2003; Siracuse, Pharm, & Sowell, 2008; Versel, 2008; Wisnicki, 2002; Wu & Wu, 2007; Zhang, 2007). For example, Wisnicki (2002) provides details of how broadband technology, an essential component of wireless technology, can be used in healthcare. While prior studies agree that wireless applications have the potential to address the endemic problems of healthcare, very limited information can be found about the determinants of such applications (Gururajan, Toleman, & Soar, 2004; Gururajan, Moloney, & Kerr, 2005). In general, the majority of the works reviewed are descriptive about the benefits of wireless handheld devices in healthcare in general, and medicine in particular. There are only a small number of studies that provide evidence-based information concerning these devices in healthcare (Fischer et al., 2003; Sax et al., 2005) (Hafeez-Baig, 2007). Furthermore, five major studies in the area of healthcare (evaluated by (Spil & Schuring, 2006) testing the Technology Acceptance Model (TAM) produced findings which were inconsistent with the body of knowledge in non-healthcare settings. With 'Perceived Ease of Use' and 'Perceived Usefulness' as the major TAM attributes, these studies found that in the health environment, 'Perceived Usefulness' is an important attribute in technology adoption, while 'Perceived Ease of Use' was found to have no effect (Spil & Schuring, 2006). This is different to findings reported in non-health IS studies, where both attributes were found to be reliable technology adoption predictors. Therefore, further empirical investigation is required to explain the reasons why this variation exists in healthcare. In addition, there is a need to explore if further attributes exist which may influence the adoption of wireless applications in the healthcare environment.

17 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/confirmatory-factor-analysis-establish-determinants/49900

Related Content

Usage of Big Data Prediction Techniques for Predictive Analysis in HIV/AIDS

Chinmayee Mohapatra, Biswaranjan Acharya, Siddhath Swarup Rautarayand Manjusha Pandey (2018). *Big Data Analytics in HIV/AIDS Research* (pp. 54-80).

www.irma-international.org/chapter/usage-of-big-data-prediction-techniques-for-predictive-analysis-in-hiv-aids/202914

Clifin: A Web-Based Geographical Information System for Time Dependant Point of Care Localization in New Brunswick, Canada

Dan Tulpan, Michelina Mancuso, Guillaume Durand, Chaouki Regouiand Luc Belliveau (2012). *International Journal of Healthcare Information Systems and Informatics* (pp. 32-47).

www.irma-international.org/article/clifin-web-based-geographical-information/75148

The Healing Hearts at Home© Mobile Application Usability and Influence on Parental Perceived Stress: A Pilot Study

Vanessa Ayer Miller, Jennifer Newcombe, Patricia Radovich, Flint Johnston, Ernesto Medina Jr.and Anna Nelson (2021). *International Journal of E-Health and Medical Communications* (pp. 90-105).

www.irma-international.org/article/the-healing-hearts-at-home-mobile-application-usability-and-influence-on-parental-perceived-stress/270905

The Power of Words: Deliberation Dialogue as a Model to Favor Patient Engagement in Chronic Care

Sarah Bigiand Giulia Lamiani (2018). *Health Care Delivery and Clinical Science: Concepts, Methodologies, Tools, and Applications* (pp. 1100-1120).

www.irma-international.org/chapter/the-power-of-words/192720

Ontological Representation and an Architecture for Homecare Pervasive Systems

Leandro Freitas, Rafael T. Pereira, Henrique G. G. Pereira, Ricardo Martini, Bruno A. Mozzaquatro, Jeferson Kasperand Giovani Librelotto (2013). *Information Systems and Technologies for Enhancing Health and Social Care* (pp. 215-234).

www.irma-international.org/chapter/ontological-representation-architecture-homecare-pervasive/75631