

## Chapter 4.6

# Planning and Control and the Use of Information Technology in Mental Healthcare Organizations

**I.J. Baars**

*Maastricht University, The Netherlands*

**G.G. Van Merode**

*Board of Maastricht University Medical Center & Maastricht University, The Netherlands*

### INTRODUCTION

Demand for mental healthcare increases. Simultaneously, the need for more patient oriented processes increases and the market develops towards more competition among providers and organizations. As a result of these developments, mental healthcare organizations are becoming more aware of efficiency and effectiveness. Often, they choose to transform to more process oriented organizations, which require changes in planning and control systems and information technology (IT). However, little is known about the required planning and control systems and IT for mental healthcare.

We argue that IT for planning and control of mental healthcare organization needs to be adaptive and support short term planning. IT has to be

adaptive to be able to support first and second order control which is needed for planning and control of mental healthcare processes. Short term planning or reactivity is needed to deal with stochasticity and variability as present in mental healthcare. These subjects are further described in the background.

This article reports the results of two studies on the use of standard care processes and IT for planning and control of mental healthcare processes. The results give insight in the needed functionalities of IT and planning and control of mental healthcare processes. The first study is a case study in a center for multidisciplinary (mental) youth care. This center implemented care programs and an automated planning tool. We studied the success of this implementation and particularly the fit between the care programs and the planning tool. In the second study we studied the characteristics

DOI: 10.4018/978-1-59904-687-7.ch021

*Table 1. Control models*

Control model condition	Unambiguous objectives	Measurable output	Known effects	Repetitive activities
Routine control	+	+	+	+
Expert control	+	+	+	-
Trial and error control	+	+	-	+
Intuitive control	+	+	-	-
Judgmental control	+	-	+/-	+/-
Political control	-	+/-	+/-	+/-

+ = condition is present

- = condition is not present

of ambulant mental healthcare processes and the actual and preferable use of planning and control models and IT.

**BACKGROUND**

Mental healthcare is often multidisciplinary and includes several professionals, disciplines, and departments within one or more organizations which all need to be planned and controlled. Consequently, the object of control is mainly professionals and patients, but also resources like rooms. We define planning as the determination of what should be done and control as the process that assures that the planned results are obtained (Van Merode, Groothuis, & Hasman, 2004).

According to Hofstede (1981), the way non-profit organizations, such as mental healthcare organizations, can be planned and controlled depends on the type of processes. The type of process can be determined by answering the following questions: is the output measurable? Are the objectives unambiguous? Are the effects of management interventions known? And, can the activities be repeated? The type of processes determines the control model and instruments (e.g., protocols, case management and budgeting) that can be applied.

Hofstede (1981) defines six different control models, as shown in Table 1. The more standard-

ized, well-defined, and structured the processes are, the more routine control can be used.

A mental healthcare organization consists of various processes. These processes are possibly different and thus need different control models. To select a control model that best suits a situation, processes have to be analyzed. Especially the distinction between routine and less routine processes is important here.

Routine processes can use standards and can be controlled by routine control or, when the activities cannot be repeated, by expert control. For routine control, and marginally for expert control and trial-and-error control, models that compare automatically what actually happens with the standards about what should happen apply. These models use feedback for control. The feedback from relevant indicators provides information that can be compared with targets. A number of care processes can be coordinated to a large extend by planning and control integrated in one system and by providing feedback from control to planning (Hofstede, 1981; Van Merode, Groothuis et al., 2004; Van Roth & Van Dierdonck, 1995). Moreover, for intuitive and judgmental and political control only vague models exist (Hofstede, 1981).

An example of control by feedback in mental healthcare is a protocol for the process of the treatment of depression. This protocol describes that the target of the indicator “number of sessions” is

6 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

[www.igi-global.com/chapter/planning-control-use-information-technology/49917](http://www.igi-global.com/chapter/planning-control-use-information-technology/49917)

## Related Content

---

### Hospital Social Media Strategies: Patient or Organization Centric?

Fay Cobb Payton and Natasha Pinto (2018). *Health Care Delivery and Clinical Science: Concepts, Methodologies, Tools, and Applications* (pp. 480-487).

[www.irma-international.org/chapter/hospital-social-media-strategies/192688](http://www.irma-international.org/chapter/hospital-social-media-strategies/192688)

### Impact of Attentional Loading and Task Constraints on Postural Control of Healthy Older Adults

Eryk Przystucha, Brontë A. Vollebregt and Carlos Zerpa (2020). *International Journal of Extreme Automation and Connectivity in Healthcare* (pp. 12-25).

[www.irma-international.org/article/impact-of-attentional-loading-and-task-constraints-on-postural-control-of-healthy-older-adults/260725](http://www.irma-international.org/article/impact-of-attentional-loading-and-task-constraints-on-postural-control-of-healthy-older-adults/260725)

### Background and Significance of Computational Convergence in Electronic Health Records

Ramneet Kaur, Renu Mishra, Jyoti Pruthi and Shruti N. Mehta (2024). *Computational Convergence and Interoperability in Electronic Health Records (EHR)* (pp. 1-26).

[www.irma-international.org/chapter/background-and-significance-of-computational-convergence-in-electronic-health-records/355568](http://www.irma-international.org/chapter/background-and-significance-of-computational-convergence-in-electronic-health-records/355568)

### Comparative Study of 4-Compartmental PK-PD Model with Effective Site Compartment for Different Parameter Set

UshaRani Shola and V Neelanarayanan (2019). *International Journal of Reliable and Quality E-Healthcare* (pp. 52-65).

[www.irma-international.org/article/comparative-study-of-4-compartmental-pk-pd-model-with-effective-site-compartment-for-different-parameter-set/219286](http://www.irma-international.org/article/comparative-study-of-4-compartmental-pk-pd-model-with-effective-site-compartment-for-different-parameter-set/219286)

### A Design Framework for a Market Intelligence System for Healthcare Sector: A Support Decision Tool in an Emergent Economy

George Jamil, Lorena Furbino, Leandro Santos, Marcus Alves, Rafael Santiago and Silvia Loyola (2013). *Handbook of Research on ICTs and Management Systems for Improving Efficiency in Healthcare and Social Care* (pp. 288-305).

[www.irma-international.org/chapter/design-framework-market-intelligence-system/78028](http://www.irma-international.org/chapter/design-framework-market-intelligence-system/78028)