

# Chapter 4.16

## Developing a User Centered Model for Ubiquitous Healthcare System Implementation: An Empirical Study

**Jongtae Yu**

*Mississippi State University, USA*

**Chengqi Guo**

*Mississippi State University, USA*

**Mincheol Kim**

*Cheju National University, South Korea*

### ABSTRACT

In the advent of pervasive computing technologies, the ubiquitous healthcare information system, or u-health system, has emerged as an innovative avenue for many healthcare management issues. Drawing upon practices in healthcare industry and conceptual developments in information systems research, this article aims to explain the latent relationships amongst user-oriented factors that lead to individual's adoption of the new technology. Specifically, this study focuses on the introduction of chronic disease u-health system. Using the ordinary line square (OLS) regression analysis, we are able to discover the insights concerning which

constructs affect service subscriber's behavioral intention of use. Based on the data collected from over 440 respondents, empirical evidences are presented to support that factors such as medical conditions, perceived need, consumer behavior, and effort expectancy significantly influence the formation of usage intention.

### INTRODUCTION

Ubiquitous computing can be defined as the "contemplation of today's computers communicating each other through wireless network in actual activities of everyday life" (Weiser, 1993a; 1993b).

Some articulate such a concept as the embedded computers in walls, refrigerator, tables, and objects in the surrounding environment (Rhodes & Mase, 2006). In other words, the computer is expected to become a ubiquitous resource, much similar to the light with a switch and water with a tap. Two functionalities, computing tasks and telecommunications, are required to realize the features of ubiquitous computing such as localized information, localized control, and resource management (Weiser, 1993a; Rhodes & Mase, 2006). The evolving mobile technology has expanded the applicability of ubiquitous computing to areas including virtual reality, head mounted display (HMD), wearable computing, and smart office room (Weiser, 1993b).

An agile, responsive, and location-aware service delivery system is highly desired by the healthcare management business. Correspondingly, ubiquitous computing allows patients to receive prompt medical care anywhere (home, office, outdoor, or hospital) and any time (24 hours/7 days), thus improving the service quality and decreasing the risk of medical treatment failures. For example, a doctor can check the status of a patient in a real-time manner using the sensor which is installed in patient's home or attached to the patient's body. In case of emergency, the sensor can detect changes of the patient's health condition in the early stage and automatically contact the designated hospital to initiate treatment procedure. In summary, we define the u-health system as the use of ubiquitous computing technologies to support expeditious and personalized communications, activities, and transactions between a medical service provider and its various stakeholders. In the literature review section, Figure 3 describes the layout of the u-health system and its associated stakeholders.

Despite the strong potential of the technology, however, the studies of ubiquitous healthcare services are not widely conducted. Existing literature has focused mostly on pure technical concerns or system development process; whereas managerial

issues and behavior perspectives of the u-health system application are largely overlooked. In this article, the authors discuss the issue of applying ubiquitous computing technology to healthcare management from end user's perspective. The main goal of this article is to identify and investigate the factors and their inter-relationships that affect end user's intention of adopting u-health system. For example, one critical factor is effort expectancy, which refers to the degree to which the user believes it is easy to use the technology. Theoretically, the effort expectancy is positively correlated with behavioral intention of use, namely, the easier the user finds to use the product, the more likely s/he will adopt it. A more thorough articulation of these factors can be found in the Research Model and Hypotheses section.

Realizing that the scope of healthcare discussion can be extremely broad, we focus our attention on chronic disease treatment in this research. A main question to be addressed is what factors influence end user's perception about the u-health system and how these factors are related with each other. Therefore, an investigation of people's subjective perception is warranted. The authors select South Korea as the place for data collection because first, people in South Korea have a high level of concern about health issues, especially chronic diseases. Largely due to the rapidly changing socio-economic structure, more than 16% of total population is reported to suffer from chronic diseases. According to a report by the Korean National Insurance Corporation (2007), South Korea is among one of the top countries that are mostly concerned with chronic diseases; second, South Korea has one of the highest mobile technology penetration rates in the world, thus providing a suitable environment for studying ubiquitous healthcare technology. In South Korea, more than 85% of Koreans use at least one cellular phone and subscribe wireless value added service (Shim, 2007).

Regarding the organization of this study, the authors first examine the current knowledge base

15 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

[www.igi-global.com/chapter/developing-user-centered-model-ubiquitous/49927](http://www.igi-global.com/chapter/developing-user-centered-model-ubiquitous/49927)

## Related Content

---

### Tele-Practice Technology: A Model for Healthcare Delivery to Underserved Populations

Thomas W. Miller, Robert D. Morgan and Jennifer A. Woods (2010). *Health Information Systems: Concepts, Methodologies, Tools, and Applications* (pp. 569-584).

[www.irma-international.org/chapter/tele-practice-technology/49887](http://www.irma-international.org/chapter/tele-practice-technology/49887)

### Factors Influencing Adoption of E-detailing as a Communication Tool: Views of Physicians of India

Saikat Banerjee and Sampada Kumar Dash (2014). *International Journal of E-Health and Medical Communications* (pp. 29-39).

[www.irma-international.org/article/factors-influencing-adoption-of-e-detailing-as-a-communication-tool/118220](http://www.irma-international.org/article/factors-influencing-adoption-of-e-detailing-as-a-communication-tool/118220)

### Demographic Differences in Telehealth Policy Outcomes

Mary Schmeida and Ramona McNeal (2010). *Health Information Systems: Concepts, Methodologies, Tools, and Applications* (pp. 1472-1480).

[www.irma-international.org/chapter/demographic-differences-telehealth-policy-outcomes/49943](http://www.irma-international.org/chapter/demographic-differences-telehealth-policy-outcomes/49943)

### An Empirical Investigation: Health Care Employee Passwords and Their Crack Times in Relationship to HIPAA Security Standards

B. Dawn Medlin and Joseph A. Cazier (2007). *International Journal of Healthcare Information Systems and Informatics* (pp. 39-48).

[www.irma-international.org/article/empirical-investigation-health-care-employee/2210](http://www.irma-international.org/article/empirical-investigation-health-care-employee/2210)

### Need of Six Sigma in Testing Laboratories: Some Explorations

Pranil Vijay Sawalakhe, Sunil V. Deshmukh and Ramesh R. Lakhe (2017). *International Journal of Reliable and Quality E-Healthcare* (pp. 24-40).

[www.irma-international.org/article/need-of-six-sigma-in-testing-laboratories/181628](http://www.irma-international.org/article/need-of-six-sigma-in-testing-laboratories/181628)