

Chapter 6

Visual Processing as Described by Contemporary Main– Stream Neuroscience

This Auxiliary Chapter complements the main part of the book by presenting the conventional views on vision for those readers who are not familiar with them. It also allows comparisons with the Holonomic Brain Theory and critiques, and also some mutual complementing of views and descriptions, although they are presently not compatible. For instance, the work of large numbers of neuroscientists provides many details and experimental data relevant for modeling purposes.

6.1 RETINA

Eye. The light enters the eye at cornea. The optic flow then proceeds through the *pupil*, a circular hole with muscle-conditioned variable radius, then through the

DOI: 10.4018/978-1-61520-785-5.ch006

lens, which bends (refracts) and focuses (because of ciliary-muscle-conditioned flexibility) the rays, to the eye's back where it impinges on the pigment-layer of *photoreceptor cells*¹ on the surface of the retina. A 2-dimensional projection of the 3-dimensional world emerges on retina.²

Retinal image. Stimulus intensities, which affect the activities of retinal cells, are mainly determined by: the light source, the surface reflectance and shape of the viewed object, the viewer's standpoint and angle of view. Each location in the image can be described by two primary characteristics of the wave-modulated stimulus (Berne & Levy, 1993):

1. *Brightness (luminance)*, the intensity of light falling on retina, which determines the photoreceptors' activation-level;
2. Spectral decomposition (frequency or *wavelength*) which determines the differential activation of different cone types, leading to the perceived color.³

Derived information. Let us mention here that other characteristics can be in later stages (up to the cortex) derived from the primary retinal features 1 and 2. Such secondary characteristics are (De Yoe & Van Essen, 1988):

3. 2-dimensional velocity of the retinal image;
4. Spatial contrast;
5. 2-dimensional orientation of the stimulus;
6. Binocular disparity (which is also a source of stereopsis, i.e. 3-dimensional vision).⁴

Sensation. Photoreceptor cells are *cones*, which are color-sensitive and used for low-sensitive (for daylight), high-acuity vision, and *rods*, which are high-sensitive (for dark), "color-blind" and provide a rough "sketch" only. In the central part of retina, *vis-a-vis* to the lens-center (on the visual axis), lies *fovea* where visual acuity is the highest, because there is extremely high density of cones. Density of rods is, in contrast, high at all other parts of the retina except at fovea and at the location where the optic nerve starts to extend towards the visual cortex. In response to a flash of light within the receptive field⁵, a photoreceptor runs a biochemical cascade process and changes voltage. Over a limited range of stimulus intensity, the change in the light-modulated signal is proportional to the change in the light stimulus (Smith in Arbib, 1995). Retina realizes a sort of scale-invariant gaussian-profile sampling (details in Van Essen & Anderson in Zornetzer, Davis & Lau, 1990, pp. 45-47).

Retinal net processing. Retina is a multi-layer neural network. Details of its non-trivial hierarchical feed-forward⁶ processing, in spite of its importance for further visual processing, will not be presented here. However, description of

46 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/chapter/visual-processing-described-contemporary-main/50505

Related Content

High-Accuracy Characterization of Ambulatory Holter Electrocardiogram Events: A Comparative Study Between Walsh-Hadamard Transform, First-Derivative-Based and Intelligent Techniques

Mohammad Reza Homaeinezhad, Seyyed Amir Hoseini Sabzevari, Ali Ghaffariand Mohammad Daevaeiha (2012). *International Journal of Systems Biology and Biomedical Technologies* (pp. 40-71).

www.irma-international.org/article/high-accuracy-characterization-ambulatory-holter/70017

Connecting the Bench with the Bedside: Translating Life Science Discoveries to Disease Treatments and Vice Versa

Yue Wang Webster, Ernst R. Dowand Mathew J. Palakal (2012). *Systemic Approaches in Bioinformatics and Computational Systems Biology: Recent Advances* (pp. 249-267).

www.irma-international.org/chapter/connecting-bench-bedside/60836

Performance Assessment of Learning Algorithms on Multi-Domain Data Sets

Amit Kumarand Bikash Kanti Sarkar (2018). *International Journal of Knowledge Discovery in Bioinformatics* (pp. 27-41).

www.irma-international.org/article/performance-assessment-of-learning-algorithms-on-multi-domain-data-sets/202362

Perspective Wall Technique for Visualizing and Interpreting Medical Data

Hela Ltifi, Mounir Ben Ayed, Ghada Trabelsiand Adel M. Alimi (2012). *International Journal of Knowledge Discovery in Bioinformatics* (pp. 45-61).

www.irma-international.org/article/perspective-wall-technique-visualizing-interpreting/77930

Evaluating a Genetics Concept Inventory

Felicia Zhangand Brett Andrew Lidbury (2013). *Bioinformatics: Concepts, Methodologies, Tools, and Applications* (pp. 29-41).

www.irma-international.org/chapter/evaluating-genetics-concept-inventory/76055