

Chapter 4.1

A Composite Indicator to Measure Agricultural Sustainability: Alternative Approaches

José A. Gómez-Limón

Institute of Agricultural Research and Training (IFAPA), Spain

Laura Riesgo

Pablo de Olavide University, Spain

ABSTRACT

This chapter deals with a comparative analysis of alternative methods of constructing composite indicators to measure the global sustainability of the agricultural sector. The comparison is implemented empirically on the irrigated agriculture of the Duero basin in Spain as a case study. For this purpose, the study employs a dataset of indicators previously calculated for different farm types and policy scenarios. The results enable us to establish a hierarchy of policy scenarios on the basis of the level of sustainability achieved. By analysing the heterogeneity of different farms types in each scenario, it is also possible to determine the main features of the most sustainable

farms in each case. All this information is useful for the support of agricultural policy design and its implementation, as we attempt to improve the sustainability of this sector.

INTRODUCTION

Agricultural sustainability does not have a single meaning. Hansen (1996) identified two broad interpretations of the concept of agricultural sustainability. The first focuses on a normative approach in response to concerns about the negative impacts of “conventional” agriculture. This approach relies on the implementation of “alternative” agriculture (ecological agriculture, conservative agriculture, etc.), as an ideological option to achieve a set of

DOI: 10.4018/978-1-60960-472-1.ch401

values that should characterize this sector. The second meaning takes a positive approach, and focuses on the ability of agricultural systems to satisfy different demands through time. As has been pointed out by Hansen (1996), only the latter meaning is useful from a scientific point of view. In this paper, therefore, we adopt this approach.

However, it is worth pointing out that the selected concept of sustainability has several difficulties that limit its empirical use in the real world. First, we have to deal with the temporal nature of sustainability. Indeed, this meaning of sustainability, which is related to the preservation of production capacity, has little practical value because of the infeasibility of long-term experiments. Second, we have to deal with the difficulty of identifying the demands that must be satisfied by the agricultural sector if it is to be regarded as sustainable. In this way, sustainability can be interpreted as a social concept that can be modified in response to the requirements of society. Thus, sustainability concept must be regarded as being both local and time-specific. Both difficulties have limited the usefulness of this concept as a criterion for guiding agricultural development.

In order to avoid the difficulties mentioned above, a wide consensus has emerged, which considers that the concept of sustainability embodies three main dimensions: environmental, economic and social (Yunlong and Smit, 1994). It can thus be assumed that an agricultural system is sustainable when the trade-offs between the objectives considered for public evaluation of its performance (economic objectives, such as income growth or macroeconomic stability; social objectives, such as equity or the cover of basic needs; and ecological objectives, such as ecosystem protection or natural resources regeneration—) reach acceptable values for society as a whole (Hediger, 1999; Stoorvogel et al., 2004). This approximation to agricultural sustainability allows its use as an operational criterion, by using a set of indicators that will cover the three dimensions mentioned above.

However, the quantification of agricultural sustainability through a set of indicators still has certain shortcomings. The main inconvenience comes from the difficulty of interpreting the complete set of indicators. In order to avoid this problem, it has been suggested that the analysis of agricultural sustainability can be tackled by aggregating this multidimensional set of indicators into a single index or composite indicator. This approach has been used, for example, by Stockle et al. (1994), Andreoli and Tellarini (2000), Pirazzoli and Castellini (2000), Sands and Podmore (2000), Rigby et al. (2001), van Calker et al. (2006) and Qiu et al. (2007). Nevertheless, the aggregation of indicators has been frequently criticised for: a) the subjectivity of the methods employed (the choice of functional forms for aggregation and weighting for individual indicators), and b) the compensability usually regarded as aggregating the individual dimensions or attributes of sustainability (additive aggregation approaches), in spite of their theoretical incommensurability. For further details see Hansen (1996), Bockstaller et al. (1997), Morse et al. (2001), Díaz-Balteiro and Romero (2004), Ebert and Welsch (2004), Munda (2005) or Böhringer and Jochem (2007).

Within this general framework, this paper has a double objective. First, from a theoretical perspective, we analyse the pros and cons of alternative methods of building composite indicators of agricultural sustainability. This is done empirically by implementing these methods in a real-world case study. Specifically, we apply these methods to quantify the global sustainability of irrigated agriculture in the Duero river basin in Spain, using an existing dataset of indicators (Riesgo and Gómez-Limón, 2005 and 2006), which covers the three dimensions of sustainability mentioned above. This set of indicators has been calculated for different farm types and future policy scenarios. This feature of the data has enabled us to consider a second objective: to analyse the real possibilities of using the concept of sustainability as a tool to guide the public management of agriculture. The

25 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/composite-indicator-measure-agricultural-sustainability/51720

Related Content

Sustainable Solid Waste Management in Developing Countries: Tanzania Outlook

Proscovia Paschal Kamugisha and Sebastian Faustin Mhanga (2020). *Sustainable Waste Management Challenges in Developing Countries* (pp. 382-408).

www.irma-international.org/chapter/sustainable-solid-waste-management-in-developing-countries/240085

Predicting Woody Plant Diversity as Key Component of Ecosystems: A Case Study in Central Greece

Alexandra D. Solomou and Athanassios Sfougaris (2019). *International Journal of Agricultural and Environmental Information Systems* (pp. 1-20).

www.irma-international.org/article/predicting-woody-plant-diversity-as-key-component-of-ecosystems/216449

A Generalization of the Orthogonal Regression Technique for Life Cycle Inventory

Antonino Marvuglia, Maurizio Cellura and Marcello Pucci (2012). *International Journal of Agricultural and Environmental Information Systems* (pp. 51-71).

www.irma-international.org/article/generalization-orthogonal-regression-technique-life/62066

Reconstruction of Missing Hourly Precipitation Data to Increase Training Data Set for ANN

Hema Nagaraja, Krishna Kant and K. Rajalakshmi (2018). *International Journal of Agricultural and Environmental Information Systems* (pp. 62-84).

www.irma-international.org/article/reconstruction-of-missing-hourly-precipitation-data-to-increase-training-data-set-for-ann/192195

Modeling Species Distribution

Yongyut Trisurat and Albertus G. Toxopeus (2011). *Land Use, Climate Change and Biodiversity Modeling: Perspectives and Applications* (pp. 171-197).

www.irma-international.org/chapter/modeling-species-distribution/53752