Chapter 6.14 Diagnostic Cost Reduction Using Artificial Neural Networks: The Case of Pulmonary Embolism

Steven Walczak University of Colorado at Denver, USA

Bradley B. Brimhall University of New Mexico, USA

Jerry B. Lefkowitz Weill Cornell College of Medicine, USA

ABSTRACT

Patients face a multitude of diseases, trauma, and related medical problems that are difficult and costly to diagnose with respect to direct costs, including pulmonary embolism (PE). Advanced decision-making tools such as artificial neural networks (ANNs) improve diagnostic capabilities for these problematic medical conditions. The research in this chapter develops a backpropagation trained ANN diagnostic model to predict the occurrence of PE. Laboratory database values for 292 patients who were determined to be at risk for a PE, with 15% suffering a confirmed PE, are collected and used to evaluate various ANN models' performance. Results indicate that using ANN diagnostic models enables the leveraging of knowledge gained from standard clinical laboratory tests, significantly improving both overall positive predictive and negative predictive performance.

DOI: 10.4018/978-1-60960-561-2.ch614

INTRODUCTION

Medical and surgical patients today face a variety of conditions that are both difficult and costly to diagnose and treat. With ever skyrocketing medical costs (Benko, 2004), the use of information technology is seen as a much-needed means to help control and potentially reduce medical direct costs (Intille, 2004). Deep vein thrombosis (DVT) and pulmonary embolism (PE) are medical conditions that are particularly difficult to diagnose in the acute setting (Mountain, 2003). Frequent usage of costly clinical laboratory tests to screen patients for further treatment is commonplace. All too commonly hospitals provide treatment to patients without PE as a preventative measure (Mountain, 2003). Furthermore, patient mortality, morbidity, and both direct and indirect costs for delayed diagnosis of these conditions may also be substantial. Recent studies show that 40 to 80 percent of patients that die from a PE are undiagnosed as having a potential PE (Mesquita et al., 1999; Morpurgo et al., 1998).

DVT may occur as the result of patient genetic and environmental factors or as a side effect of lower extremity immobility (e.g., following surgery). When a blood clot in the veins of a lower extremity breaks away, it may travel to the lungs and lodge in the pulmonary arterial circulation causing PE. If the clot is large enough it may wedge itself into the large pulmonary arteries leading to an acute medical emergency with a significant mortality rate. Approximately 2 million people annually experience DVT, with approximately 600,000 developing a PE and approximately 10% of these PE episodes result in mortality (Mesquita et al. 1999; Labarere et al., 2004). Documented occurrence of DVT in postoperative surgical populations ranges from 10% (Hardwick & Colwell, 2004) to 28% (Blattler et al., 2004).

Direct costs associated with DVT and PE come from the expensive diagnostic and even more expensive treatment protocols. It may be possible to lower these direct costs, especially when additional testing or treatment may be ruled out due to available knowledge. Artificial neural network (ANN) systems enable the economic examination (Walczak, 2001) and nonlinear combination of various readily available clinical laboratory tests. Laboratory tests typically performed on surgical patients, for example, blood chemistry, form the foundation for analysis and diagnostic model development.

One such test is the D-dimer assay that measures patient plasma for the concentration of one molecular product released from blood clots. When blood vessels are injured or when the movement of blood is too slow through veins of lower extremities, blood may begin to clot by initiating a series of steps in which fibrin molecules are crossed linked by thrombin to form a structure that entraps platelets and other coagulation molecules-a blood clot. As healing begins to occur, plasmin begins to break down the clot and releases, among other things, D-dimer molecules. D dimers are actually small fragments of cross-linked fibrin and provide the basis for assessing blood clotting activity. Patients with DVT and PE frequently have elevated levels of D-dimer in their plasma. Consequently, many hospitals now employ the D-dimer assay as a first test in the diagnostic pathway for these conditions. Usage of the results of a D-dimer assay effectively reduces the direct costs of DVT by reducing the requirement for downstream testing and treatment, specifically Doppler ultrasound tests (Wells, 2003). The use of nonparametric models enables the analysis of laboratory tests without regard to the population distribution, which may be a problematic factor when combining more than one laboratory test for the diagnostic prediction. Additionally, ANNs provide nonlinear modeling capabilities that may be beneficial in combining pathology tests.

The research reported in this chapter will examine the efficacy of using ANN models to predict the likelihood of a PE in surgical patient populations. A corollary research question is whether less invasive and less costly diagnostic 16 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage: www.igi-global.com/chapter/diagnostic-cost-reduction-using-artificial/53683

Related Content

The Implementation of Innovative Technologies in Healthcare

Eddy M.M. Adang (2011). *Clinical Technologies: Concepts, Methodologies, Tools and Applications (pp. 1-12).*

www.irma-international.org/chapter/implementation-innovative-technologies-healthcare/53573

E-Medical Education

D. John Doyle (2011). Clinical Technologies: Concepts, Methodologies, Tools and Applications (pp. 190-209).

www.irma-international.org/chapter/medical-education/53585

Women's Health Informatics in the Primary Care Setting

Gareth Parry (2011). Clinical Technologies: Concepts, Methodologies, Tools and Applications (pp. 1518-1529).

www.irma-international.org/chapter/women-health-informatics-primary-care/53663

Predicting Ambulance Diverson

Abey Kuruvillaand Suraj M. Alexander (2011). *Clinical Technologies: Concepts, Methodologies, Tools and Applications (pp. 1393-1402).*

www.irma-international.org/chapter/predicting-ambulance-diverson/53655

Proliferation and Regeneration: Methodologies in Cancer Treatment and Post-Treatment Tissue Reconstruction

George I. Lambrou, Maria Adamakiand Apostolos Zaravinos (2013). *Medical Advancements in Aging and Regenerative Technologies: Clinical Tools and Applications (pp. 31-52).* www.irma-international.org/chapter/proliferation-regeneration-methodologies-cancer-treatment/71975