

Chapter 3

A Comparison of Principal Component–Based and Multivariate Regression of Cardiac Disease

Fox Underwood

University of Calgary, Canada

Stefania Bertazzon

University of Calgary, Canada

ABSTRACT

Selecting factors suitable to use in a regression model is often a complicated process: the researcher strives to retain all theoretically important factors while avoiding high correlations among independent variables. This chapter models cardiac disease and compares the explanatory ability of component-based multivariate regression models, created through the use of principal component analysis (PCA), with that of direct variable-based, multivariate regression models. The variable-based demographic and socio-economic model contains education, sex, and 3 age factors; in contrast, the component-based model contains age as well as several modifiable risk factors: education, income, family, and housing factors. Moreover, the latter model also has statistically higher explanatory power. Components made through data reduction techniques may not always be interpretable, but, given closer examination of individual components, a component-based model becomes more interpretable. Further, all important factors will potentially be present in models. As such, component-based modelling can be a useful tool for research and public health planning. A key limitation of this work, to be addressed in future research, is the use of a variable (cardiac catheterisation procedures) that remains a crude proxy for cardiovascular disease. More effective analysis will be performed as data becomes available. Exploration into the relationship of factor and their spatial patterns will also be considered.

DOI: 10.4018/978-1-4666-1924-1.ch003

INTRODUCTION

Many primary health concerns of contemporary societies are inherently spatial: effective accessibility to health care services, prompt and efficient response to epidemic outbreaks, detection and monitoring of environmental health hazards, and consequent urban and regional planning. Management and planning decisions are often supported by quantitative models; indeed the use of quantitative and statistical methods in health research is well established, and the application of spatial analytical methods has gained acceptance over the last several years (Elliott et al., 2000; Waller and Gotway, 2003; Elliott and Wartenberg, 2004). It has been argued that the integration of analytical and visual methods can improve the effectiveness of spatial analysis as a decision support tool for policy and management (Guo, 2007). Further, it has been argued that, though some difficulties still exist, geographic information science (GIS) has a potential role in improving public health (Rushton, 2000).

Despite this potential in assisting management and planning decisions, the use of quantitative methods to spatial data remains a delicate process, and its simplistic application does not necessarily yield reliable and relevant results. One well known problem is that of estimating uncertainty, which stems from two intrinsic properties of geographical phenomena: spatial dependence (i.e. near things are more related than distant things) and spatial non-stationarity (i.e. inconstant variability over space) (Cliff and Ord, 1981). Violation of either assumption inflates the variance—and hence the uncertainty—of the regression estimates, resulting in less reliable models (Anselin, 1988). An additional source of model instability is cross-correlation of the explanatory variables, known as multicollinearity in the model (McGarigal et al., 2001). A common strategy to avoid multicollinearity or near-multicollinearity is a drastic model selection, where only explanatory variables that display low cross-correlation are retained in the model (McGarigal et al., 2001).

This paper presents the use of multivariate regression analysis to identify demographic and socio-economic variables significantly associated with cardiovascular disease prevalence in a Calgary: a large Canadian city. In consideration of the available data (i.e. spatially aggregated records) spatial regression techniques are applied in the presence of spatial dependence to enhance the reliability of the model parameters. In response to multicollinearity, the use of principal component analysis (PCA) (McGarigal et al., 2001) is proposed, leading to the comparison of regression models which make use of orthogonal components and of selected uncorrelated variables, respectively.

Selecting factors suitable to use in a regression model is often a complicated process: the researcher strives to retain all theoretically important factors while avoiding high correlations among independent variables. This tends to result in a compromise rather than a success as not all important factors can always be retained. ‘Rules of thumb’ and intuition are also often used when deciding on factors to retain; the larger the pool of factors, the more time-consuming this process is. Such effort can be reduced by employing sets of components gleaned through data reduction techniques rather than sets of variables.

BACKGROUND

Study Area and Data

Cardiovascular diseases are among the leading causes of death in the developed world (Kaplan and Keil, 1993; Manuel et al., 2003; Gerber et al. 2006; Canadian Heart Health Strategy and Action Plan, 2009). Disease occurrence is related to personal characteristics such as age, sex, genetic background, and the simultaneous presence of other conditions; all these are known as non-modifiable risk factors. In addition to these, disease prevalence has been found in association with a number of modifiable risk factors,

16 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/comparison-principal-component-based-multivariate/69047

Related Content

The Role of GIS in City Competitiveness

Khan R. Rahaman and Júlia M. Lourenço (2013). *Geographic Information Systems: Concepts, Methodologies, Tools, and Applications* (pp. 1476-1501).

www.irma-international.org/chapter/role-gis-city-competitiveness/70517

Practical Approaches to Spatial Estimation of Disaster-Affected Populations

Lisa Jordan, Benjamin Watkins, Patrick Biegon, Margaret Mwangi and Rob Rose (2010). *International Journal of Applied Geospatial Research* (pp. 31-48).

www.irma-international.org/article/practical-approaches-spatial-estimation-disaster/45129

Integration of Satellite Remote Sensing Techniques and Landscape Metrics to Characterize Land Cover Change and Dynamics

Carmelo Riccardo Fichera, Giuseppe Modica and Maurizio Pollino (2013). *Geographic Information Analysis for Sustainable Development and Economic Planning: New Technologies* (pp. 228-244).

www.irma-international.org/chapter/integration-satellite-remote-sensing-techniques/69060

VGI in the Geoweb: An Experiment to Test Data Reliability

Michael Buzzelli, David Brown, Kenwoo Lee and Justin Mullan (2017). *Volunteered Geographic Information and the Future of Geospatial Data* (pp. 65-75).

www.irma-international.org/chapter/vgi-in-the-geoweb/178799

A New Spatial Transformation Scheme for Preventing Location Data Disclosure in Cloud Computing

Min Yoon, Hyeong-il Kim, Miyoung Jang and Jae-Woo Chang (2016). *Geospatial Research: Concepts, Methodologies, Tools, and Applications* (pp. 1752-1776).

www.irma-international.org/chapter/a-new-spatial-transformation-scheme-for-preventing-location-data-disclosure-in-cloud-computing/149575