Chapter 14

Can Activated Platelet Rich Plasma Combined with Adipose-Derived Stem Cells Be Used to Treat Skin Wrinkles? A Mechanism Study

Phuc Van Pham

Vietnam National University, Vietnam

Loan Thi-Tung Dang

Vietnam National University, Vietnam

Nhung Hai Truong

Vietnam National University, Vietnam

Ngoc Kim Phan

Vietnam National University, Vietnam

ABSTRACT

In recent years, Platelet Rich Plasma (PRP) and Adipose-Derived Stem Cells (ADSCs) have been used separately for many clinical applications, especially skin rejuvenation. A combined injection of PRP and ADSCs could therefore be used to treat skin wrinkles. However, there are controversies and reports with conflicting results regarding the efficacy of this treatment. The authors aimed to determine the anti-wrinkle and skin rejuvenation mechanism of combined PRP and ADSCs treatment. The effects of PRP and ADSCs isolated from the same consenting donors were evaluated using in vitro and in vivo models. The in vitro effects of PRP and ADSCs on dermal fibroblast proliferation, collagen production, and inhibition of Matrix Metalloproteinase-1 (MMP-1) production were investigated using a co-culture model. Fibroblasts and ADSCs were cultured within the same dish, but in two separate cavities (using an insert plate), in the presence of the same PRP-supplemented medium. In vivo, the authors evaluated the effects of combined PRP and ADSCs on skin histochemistry, including changes in the dermal layer and collagen production in photo-aged skin (mice). They also determined the survival and differentiation of grafted ADSCs. The results show that combined PRP and ADSCs strongly stimulate in vitro fibroblast proliferation, collagen production, and inhibition of MMP-1 synthesis. Intra-dermal co-injection of PRP and ADSCs was observed to stimulate increased dermal layer thickness and collagen production compared with the untreated group. These results indicate that a combined PRP and ADSC injection can reduce wrinkles more effectively than either PRP or ADSC alone, and provide insight into the clinical use of PRP combined with ADSCs for dermal applications, particularly skin rejuvenation.

DOI: 10.4018/978-1-4666-2506-8.ch014

INTRODUCTION

Skin wrinkles are the most obvious sign of ageing skin, usually characterized by small and fine wrinkles, roughness, dryness, laxity, and changes in pigmentation. This phenomenon is due to epidermal thinning and collagen degradation, a process effected and enhanced by ultraviolet rays, so is also referred to as "photo-aged" (Chung, 2001, 2003). Following skin collagen degradation, elasticity is reduced and wrinkles are formed (Fisher, 2008). Wrinkles usually appear on the face skin after the age of 28. The inhibition of skin ageing and therefore wrinkles is of great interest to a majority of people.

Previously, skin wrinkles have been treated using a range of therapies, including collagen, hyaluronic acid, and adipose injections; however, results have been disappointing. Recently, PRP and ADSCs have been used widely across many clinical fields, especially for skincare and cosmetic surgery. PRP is an enriched plasma, obtained by a simple centrifugation technique. It has been reported to stimulate bone adhesion and wound healing (Froum, 2002; Marx, 1998; Petrungaro, 2001; Robiony, 2002; Margolis, 2001; Welsh, 2000; Man, 2001; Bhanot, 2002). PRP contains a pool of at least seven Growth Factors (GFs), including Epidermal Growth Factor (EGF), Platelet Derived Growth Factor (PDGF), Transforming Growth Factor Beta (TGF-β), Vascular Endothelial Growth Factor (VEGF), Fibroblast Growth Factor (FGF), Insulin-like Growth Factor (IGF), and Keratinocyte Growth Factor (KGF). The therapeutic influence of PRP is due to the high concentration of these growth factors compared with normal plasma. Many of these GFs have important roles in wound healing and tissue regeneration. EGF can stimulate proliferation, differentiation, and survival of cells (Herbst, 2004). KGF, also known as FGF7, is a GF present in the epithelialization phase of wound healing. It stimulates keratinocyte proliferation to cover the wound and form the

epithelium. These GFs are reported to improve photo-aged skin, enhance collagen production, and stimulate keratinocyte proliferation within skin (Fitzpatrick, 2003; Ehrlich, 2006; Gold, 2007). Results from previous studies have shown PRP to have many benefits for skin regeneration and anti-aging. PRP injection into nude mice resulted in rejuvenate photo-aged skin (Cho, 2011). PRP has also been shown to stimulate expression of type I collagen (col-I) and MMP-1 protein in dermal fibroblasts (Kim, 2011) and to increase expression of G1 cycle regulators, col-I, and MMP-1 to accelerate wound healing (Cho, 2012). PRP used in combination with fractional laser treatment has been successful for skin rejuvenation. This combination can stimulate keratinocyte and fibroblast proliferation and collagen production, so increase dermal elasticity.

Recently, Adipose-Derived Stem Cells (ADSCs) were used for skin rejuvenation and anti-aging applications. ADSCs are multipotent stem cells derived from adipose tissue (Zuk, 2002; Bunnell, 2008; Witkowska-Zimny, 2011; Yu, 2011). ADSCs can differentiate into various lineages, including adipocytes (Chen, 2012); insulin producing cells (Kim, 2010), endothelial cells (Colazzo, 2010; Colazzo, 2010; Zhang, 2011; Marino, 2012); osteoblasts (Xu, 2005; Safwani, 2011), hepatocytes (Al Battah, 2011; Snykers, 2011); chondrocytes (Estes, 2010; Boeuf, 2010; Musumeci, 2011), and neuronal-like cells (Gardin, 2011). ADSC transplantation has been reported to benefit skin rejuvenation by triggering dermal wound healing and inhibiting photo-aging (Kim, 2007) and also by inhibiting UVB-induced wrinkles and stimulating collagen synthesis (Kim, 2009, 2011). Furthermore, ADSCs are reported to improve autologous fat transplantation, exhibit an antioxidant action and whitening effect (Matsumoto, 2006; Kim, 2008) and increase angiogenesis (Kim, 2011). There exist two reported mechanisms explaining how ADSCs provide these therapeutic effects on skin. The first mechanism 15 more pages are available in the full version of this document, which may be purchased using the "Add to Cart" button on the publisher's webpage:

www.igi-global.com/chapter/can-activated-platelet-rich-plasma/71987

Related Content

Interactive Information Retrieval as a Step Towards Effective Knowledge Management in Healthcare

Jörg Ontrupand Helge Ritter (2005). Clinical Knowledge Management: Opportunities and Challenges (pp. 52-71).

www.irma-international.org/chapter/interactive-information-retrieval-step-towards/6577

Trust and Clinical Information Systems

Rania Shibl, Kay Fielden, Andy Bissettand Den Pain (2011). *Clinical Technologies: Concepts, Methodologies, Tools and Applications (pp. 1922-1933).*

www.irma-international.org/chapter/trust-clinical-information-systems/53689

Meta-Analysis Approach for the Identification of Molecular Networks Related to Infections of the Oral Cavity

A. Daskalakiand A. Rasche (2010). *Informatics in Oral Medicine: Advanced Techniques in Clinical and Diagnostic Technologies (pp. 237-250).*

www.irma-international.org/chapter/meta-analysis-approach-identification-molecular/40448

The Impact of Information Technology across Small, Medium, and Large Hospitals

Stacy Bourgeois, Edmund Praterand Craig Slinkman (2011). *New Technologies for Advancing Healthcare and Clinical Practices (pp. 347-361).*

www.irma-international.org/chapter/impact-information-technology-across-small/55153

Case Study: Research Matchmaker, an Advanced Nursing Practice Informatics Application

Gail Heaberg (2011). Evidence-Based Practice in Nursing Informatics: Concepts and Applications (pp. 217-236).

www.irma-international.org/chapter/case-study-research-matchmaker-advanced/48934